OCR
Oxford Cambridge and RSA

Thursday 22 May 2025 – Afternoon

A Level Further Mathematics B (MEI)

Y420/01 Core Pure

Time allowed: 2 hours 40 minutes

T869069613

You must have:
- the Printed Answer Booklet
- the Formulae Booklet for Further Mathematics B (MEI)
- a scientific or graphical calculator

QP

INSTRUCTIONS
- Use black ink. You can use an HB pencil, but only for graphs and diagrams.
- Write your answer to each question in the space provided in the **Printed Answer Booklet**. If you need extra space use the lined pages at the end of the Printed Answer Booklet. The question numbers must be clearly shown.
- Fill in the boxes on the front of the Printed Answer Booklet.
- Answer **all** the questions.
- Where appropriate, your answer should be supported with working. Marks might be given for using a correct method, even if your answer is wrong.
- Give your final answers to a degree of accuracy that is appropriate to the context.
- Do **not** send this Question Paper for marking. Keep it in the centre or recycle it.

INFORMATION
- The total mark for this paper is **144**.
- The marks for each question are shown in brackets **[ ]**.
- This document has **12** pages.

ADVICE
- Read each question carefully before you start your answer.

© OCR 2025 [Y/508/5592]
DC (DE/SW) 355352/5

OCR is an exempt Charity

stemup.co.uk

Section A (33 marks)

1 The complex number z satisfies the equation $z + 2iz^* + 1 - 4i = 0$.

You are given that $z = x + iy$, where x and y are real numbers.

Determine the values of x and y. **[4]**

2 **In this question you must show detailed reasoning.**

Find the acute angle between the planes $2x - y + 2z = 5$ and $x + 2y + z = 8$. **[4]**

3 Using standard summation formulae, show that, for integers $n \geqslant 1$,

$1 \times 3 + 2 \times 4 + ... + n \times (n+2) = \frac{1}{6}n(n+1)(an+b)$,

where a and b are integers to be determined. **[5]**

4 **(a)** You are given that **M** and **N** are non-singular 2×2 matrices.

Write down the product rule for the inverse matrices of **M**, **N** and **MN**. **[1]**

(b) Verify this rule for the matrices **M** and **N**, where

$\mathbf{M} = \begin{pmatrix} a & 1 \\ 0 & 1 \end{pmatrix}$ and $\mathbf{N} = \begin{pmatrix} 0 & -1 \\ 1 & b \end{pmatrix}$ and a and b are non-zero constants. **[6]**


5 The cubic equation $2x^3 - 3x + 4 = 0$ has roots α, β and γ.

Determine a cubic equation with integer coefficients whose roots are $\frac{1}{2}(\alpha+1)$, $\frac{1}{2}(\beta+1)$ and $\frac{1}{2}(\gamma+1)$. **[4]**

© OCR 2025

stemup.co.uk

6 The figure below shows the curve with cartesian equation $(x^2 + y^2)^2 = xy$.



(a) Show that the polar equation of the curve is $r^2 = a \sin b\theta$, where a and b are positive constants to be determined. **[3]**

(b) Determine the exact maximum value of r. **[2]**

(c) Determine the area enclosed by **one** of the loops. **[4]**

© OCR 2025

stemup.co.uk

Section B (111 marks)

7 **In this question you must show detailed reasoning.**

By first expressing $\frac{1}{x^2-4}$ in partial fractions, show that $\int_3^\infty \frac{1}{x^2-4}\mathrm{d}x = \frac{1}{m}\ln n$, where m and n are integers to be determined. **[8]**

8 The function $\mathrm{f}(x)$ is defined as $\mathrm{f}(x) = \ln(1+x)$, for $x > -1$.

(a) Prove by mathematical induction that the nth derivative of $\mathrm{f}(x)$, $\mathrm{f}^{(n)}(x)$, for all $n \geqslant 1$, is given by $\mathrm{f}^{(n)}(x) = \frac{(-1)^{n+1}(n-1)!}{(1+x)^n}$. **[4]**


(b) Hence prove that $\ln(1+x) = x - \frac{x^2}{2} + \frac{x^3}{3} - ... + \frac{(-1)^{n+1}x^n}{n} + ...$ for $-1 < x \leqslant 1$.

[You are not required to show this series for $\ln(1+x)$ converges for $-1 < x \leqslant 1$.] **[3]**

© OCR 2025

stemup.co.uk

9 The figure below shows an Argand diagram with a **regular** pentagon ABCDE. The point A represents the real number 1. The point B represents the complex number w.



(a) (i) Write down, in terms of w, the complex numbers represented by the points C, D and E. **[1]**

(ii) Write down an equation whose roots are the complex numbers represented by the points A, B, C, D and E. **[1]**

(iii) Show that the sum of these roots is zero. **[2]**

(b) (i) Find w. Give your answer in the form $r(\cos\theta + \mathrm{i}\sin\theta)$, where $r > 0$ and $\theta = k\pi$, where k is a positive constant to be found. **[1]**

(ii) By considering the line segment AB, show that the length of each side of the pentagon is $2\sin\frac{\pi}{5}$. **[5]**

10 In this question you must show detailed reasoning.

Evaluate $\int_0^{\frac{1}{2}} \frac{2}{x^2 - x + 1}\,\mathrm{d}x$. Give your answer in exact form. **[4]**

stemup.co.uk

11 The lines l_1 and l_2 have equations

$$l_1: \frac{x-3}{a} = \frac{y+1}{b} = \frac{z-2}{1} \qquad l_2: \mathbf{r} = 2\mathbf{i} + \mathbf{j} + 3\mathbf{k} + \lambda(-\mathbf{i} + c\mathbf{j} + 2\mathbf{k})$$

where a, b and c are constants.

(a) In the case where $c = 0$ and l_1 and l_2 intersect at right angles, determine the coordinates of the point of intersection of the two lines. **[6]**

(b) Now consider the case where $c = 4$ and lines l_1 and l_2 are parallel.

(i) Find the values of a and b. **[3]**

(ii) Determine the distance between the two lines. **[5]**

12 In this question you must show detailed reasoning.

The roots of the equation $z^4 - z^3 + cz^2 + dz + 18 = 0$ are α, $\frac{2}{\alpha}$, β and $-\beta$.

Determine, in any order, the exact values of the following.

- The **four** roots of the equation
- The value of c
- The value of d **[8]**

13 The gradient of a curve $y = \mathrm{f}(x)$ satisfies the differential equation $x\frac{\mathrm{d}y}{\mathrm{d}x} - 2y = 2 + x^2$.

(a) Show that the integrating factor for this differential equation is x^{-2}. **[3]**

(b) You are given that the curve passes through the point (1, 0).

By solving this differential equation, determine the exact x-coordinate of the stationary point on the curve $y = \mathrm{f}(x)$. **[8]**

© OCR 2025

stemup.co.uk

14 **(a)** By using the definition of $\cosh x$ and $\sinh x$ in terms of e^x and e^{-x}, show that $\cosh^2 x + \sinh^2 x \equiv \cosh 2x$. **[2]**

(b) The transformation T of the plane has associated matrix **M**, where $\mathbf{M} = \begin{pmatrix} \cosh x & \sinh x \\ \sinh x & \cosh x \end{pmatrix}$ and $x > 0$.

Show that T transforms the unit square with coordinates (0, 0), (1, 0), (0, 1) and (1, 1) to a rhombus of unit area. **[6]**

(c) You are given that the length of each side of the rhombus is 2 units.

Determine the exact value of x. Give your answer in logarithmic form. **[2]**

15 **(a)** Show that $(3 - e^{4i\theta})(3 - e^{-4i\theta}) = a + b\cos 4\theta$, where a and b are integers to be determined. **[2]**

The infinite series C and S are defined as follows.

$$C = \cos\theta + \frac{1}{3}\cos 5\theta + \frac{1}{9}\cos 9\theta + \frac{1}{27}\cos 13\theta + ...$$

$$S = \sin\theta + \frac{1}{3}\sin 5\theta + \frac{1}{9}\sin 9\theta + \frac{1}{27}\sin 13\theta + ...$$

(b) Show that $C + iS = \dfrac{3e^{i\theta}}{3 - e^{4i\theta}}$. **[4]**


(c) Hence show that $C = \dfrac{9\cos\theta - 3\cos 3\theta}{10 - 6\cos 4\theta}$. **[3]**

© OCR 2025

stemup.co.uk

16 In this question you must show detailed reasoning.

The diagram shows the curve with equation $y = \dfrac{x+3}{\sqrt{x^2+9}}$.



The region R, shown shaded in the diagram, is bounded by the curve, the x-axis, the y-axis, and the line $x = 4$.

(a) Determine the area of R. Give your answer in the form $p + \ln q$ where p and q are integers to be determined. **[6]**

The region R is rotated through 2π radians about the x-axis.

(b) Determine the volume of the solid of revolution formed. Give your answer in the form $\pi\left(a + b\ln\left(\dfrac{c}{d}\right)\right)$ where a, b, c and d are integers to be determined. **[6]**

© OCR 2025

stemup.co.uk

17 A researcher is modelling the height of a particular type of tree over its lifetime.

Data suggests that the maximum possible height of this type of tree over its lifetime is double the height of the tree 5 years after planting.

It is given that, t years after planting a seed for this type of tree, the corresponding height of the tree is h m, and that $h = 0$ when $t = 0$.

(a) The researcher first models the height of the tree by assuming that the rate of increase of h is proportional to $(20 - h)$, with constant of proportionality 0.2.

(i) Write down the first order differential equation for this model. **[1]**

(ii) Show that this model predicts that the maximum possible height of the tree is 20 m. **[1]**

(iii) Show by integration that $h = 20(1 - e^{-0.2t})$. **[4]**

(iv) Determine whether this model's prediction for the height of the tree 5 years after planting is consistent with the maximum possible height of the tree being 20 m. **[2]**

(b) The researcher refines the model for the height of the tree using the following second order differential equation.

$$\frac{d^2h}{dt^2} + 0.3\frac{dh}{dt} + 0.02h = 0.4$$

(i) Determine the general solution of this second order differential equation. **[4]**

(ii) Show that the refined model also predicts that the maximum possible height of the tree is 20 m. **[1]**

Further research determines that the initial rate of growth of this type of tree is 2.9 metres per year.

(iii) By applying the initial conditions to find the particular solution of this differential equation, determine whether the refined model's prediction for the height of the tree 5 years after planting is consistent with the maximum possible height of the tree being 20 m. **[5]**

END OF QUESTION PAPER

© OCR 2025

stemup.co.uk

BLANK PAGE

© OCR 2025

stemup.co.uk

BLANK PAGE

© OCR 2025

stemup.co.uk

OCR
Oxford Cambridge and RSA

Copyright Information

OCR is committed to seeking permission to reproduce all third-party content that it uses in its assessment materials. OCR has attempted to identify and contact all copyright holders whose work is used in this paper. To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced in the OCR Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download from our public website (www.ocr.org.uk) after the live examination series.

If OCR has unwittingly failed to correctly acknowledge or clear any third-party content in this assessment material, OCR will be happy to correct its mistake at the earliest possible opportunity.

For queries or further information please contact The OCR Copyright Team, The Triangle Building, Shaftesbury Road, Cambridge CB2 8EA.

OCR is part of Cambridge University Press & Assessment, which is itself a department of the University of Cambridge.

© OCR 2025

stemup.co.uk

OCR
Oxford Cambridge and RSA

Thursday 22 May 2025 – Afternoon

A Level Further Mathematics B (MEI)

Y420/01 Core Pure

Printed Answer Booklet

Time allowed: 2 hours 40 minutes

AB

You must have:
- Question Paper Y420/01 (inside this document)
- the Formulae Booklet for Further Mathematics B (MEI)
- a scientific or graphical calculator

Please write clearly in black ink. **Do not write in the barcodes.**

Centre number | | | | | | Candidate number | | | | |

First name(s) StemUp Model Solution

Last name

INSTRUCTIONS
- Use black ink. You can use an HB pencil, but only for graphs and diagrams.
- Write your answer to each question in the space provided in the **Printed Answer Booklet**. If you need extra space use the lined pages at the end of the Printed Answer Booklet. The question numbers must be clearly shown.
- Answer **all** the questions.
- Where appropriate, your answer should be supported with working. Marks might be given for using a correct method, even if your answer is wrong.
- Give your final answers to a degree of accuracy that is appropriate to the context.

INFORMATION
- This document has **24** pages.

ADVICE
- Read each question carefully before you start your answer.

© OCR 2025 [Y/508/5592]
DC (DE) 355356/2
OCR is an exempt Charity

stemup.co.uk

Section A (33 marks)

1

$z^* = x - iy$

$x + iy + 2i(x - iy) + 1 - 4i = 0$

$x + iy + 2ix - 2i^2y + 1 - 4i = 0$

$x + iy + 2ix + 2y + 1 - 4i = 0$ (since $i^2 = -1$)

Real parts: $x + 2y + 1 = 0$

Imaginary parts: $y + 2x - 4 = 0 \Rightarrow y = 4 - 2x$

$x + 2(4 - 2x) + 1 = 0$

$-3x + 9 = 0 \Rightarrow x = 3$

$y = 4 - 2(3) = -2$

$x = 3, \quad y = -2$

2

Normal vectors: $n_1 = 2i - j + 2k, \quad n_2 = i + 2j + k$

$n_1 \cdot n_2 = 2(1) + (-1)(2) + 2(1) = 2 - 2 + 2 = 2$

$|n_1| = \sqrt{2^2 + (-1)^2 + 2^2} = \sqrt{9} = 3$

$|n_2| = \sqrt{1^2 + 2^2 + 1^2} = \sqrt{6}$

$\cos\theta = (n_1 \cdot n_2) / (|n_1||n_2|)$

$\cos\theta = \frac{2}{3\sqrt{6}} = 0.27216...$

$\theta = \cos^{-1}(0.27216...) = 74.207...°$

This is acute, so the acute angle between the planes is 74.2° (3 s.f.)

© OCR 2025

3 $1 \times 3 + 2 \times 4 + \ldots + n(n+2) = \sum_{r=1}^{n} r(r+2)$

$= \sum_{r=1}^{n} r^2 + 2\sum_{r=1}^{n} r$

$= \frac{1}{6}n(n+1)(2n+1) + 2 \times \frac{1}{2}n(n+1)$

$= \frac{1}{6}n(n+1)(2n+1) + \frac{6}{6}n(n+1)$

$= \frac{1}{6}n(n+1)[(2n+1)+6]$

$= \frac{1}{6}n(n+1)(2n+7)$

So $a = 2$ and $b = 7$

© OCR 2025

stemup.co.uk

4(a)

$(MN)^{-1} = N^{-1}M^{-1}$

4(b)

$\det M = a(1) - 1(0) = a$

$M^{-1} = \frac{1}{a}\begin{pmatrix} 1 & -1 \\ 0 & a \end{pmatrix}$

$\det N = 0(b) - (-1)(1) = 1$

$N^{-1} = \frac{1}{1}\begin{pmatrix} b & 1 \\ -1 & 0 \end{pmatrix} = \begin{pmatrix} b & 1 \\ -1 & 0 \end{pmatrix}$

$MN = \begin{pmatrix} a & 1 \\ 0 & 1 \end{pmatrix}\begin{pmatrix} 0 & -1 \\ 1 & b \end{pmatrix} = \begin{pmatrix} 0+1 & -a+b \\ 0+1 & 0+b \end{pmatrix} = \begin{pmatrix} 1 & b-a \\ 1 & b \end{pmatrix}$

$\det(MN) = 1(b) - (b-a)(1) = a$

$(MN)^{-1} = \frac{1}{a}\begin{pmatrix} b & a-b \\ -1 & 1 \end{pmatrix}$

$N^{-1}M^{-1} = \begin{pmatrix} b & 1 \\ -1 & 0 \end{pmatrix} \times \frac{1}{a}\begin{pmatrix} 1 & -1 \\ 0 & a \end{pmatrix}$

$= \frac{1}{a}\begin{pmatrix} b+0 & -b+a \\ -1+0 & 1+0 \end{pmatrix} = \frac{1}{a}\begin{pmatrix} b & a-b \\ -1 & 1 \end{pmatrix}$

So $(MN)^{-1} = N^{-1}M^{-1}$, so the rule is verified.

© OCR 2025

5 Let $y = \frac{1}{2}(x + 1)$, so $x = 2y - 1$

Substitute into $2x^3 - 3x + 4 = 0$:

$2(2y - 1)^3 - 3(2y - 1) + 4 = 0$

$(2y - 1)^3 = 8y^3 - 12y^2 + 6y - 1$

$2(8y^3 - 12y^2 + 6y - 1) - 6y + 3 + 4 = 0$

$16y^3 - 24y^2 + 12y - 2 - 6y + 7 = 0$

$16y^3 - 24y^2 + 6y + 5 = 0$

So a required equation is $16x^3 - 24x^2 + 6x + 5 = 0$

© OCR 2025

stemup.co.uk

6(a)

$x = r\cos\theta, \quad y = r\sin\theta, \quad x^2 + y^2 = r^2$

$(r^2)^2 = (r\cos\theta)(r\sin\theta)$

$r^4 = r^2\sin\theta\cos\theta$

$r^2 = \sin\theta\cos\theta = \frac{1}{2}\sin 2\theta$

So $r^2 = \frac{1}{2}\sin 2\theta, \quad a = \frac{1}{2}, \quad b = 2$

6(b)

r^2 is greatest when $\sin 2\theta = 1$, i.e. $\theta = \frac{\pi}{4}$

Maximum $r^2 = \frac{1}{2}$, so maximum $r = \sqrt{1/2}$

Maximum value of $r = \frac{1}{\sqrt{2}} \quad \left(= \frac{\sqrt{2}}{2}\right)$

6(c)

One loop is traced for $0 \leqslant \theta \leqslant \frac{\pi}{2}$ (where $\sin 2\theta \geqslant 0$)

Area $= \frac{1}{2}\int_0^{\pi/2} r^2\, d\theta = \frac{1}{2}\int_0^{\pi/2} \frac{1}{2}\sin 2\theta\, d\theta$

$= \frac{1}{4}\int_0^{\pi/2} \sin 2\theta\, d\theta$

$= \frac{1}{4}\left[-\frac{1}{2}\cos 2\theta\right]_0^{\pi/2}$

$= -\frac{1}{8}(\cos\pi - \cos 0)$

$= -\frac{1}{8}(-1 - 1)$

$= \frac{1}{4}$

Area enclosed by one loop $= \frac{1}{4}$

© OCR 2025

Section B (111 marks)

7

$\frac{1}{x^2-4} = \frac{1}{(x-2)(x+2)} = \frac{A}{x-2} + \frac{B}{x+2}$

$1 \equiv A(x+2) + B(x-2)$

$x = 2$: $1 = 4A \Rightarrow A = \frac{1}{4}$

$x = -2$: $1 = -4B \Rightarrow B = -\frac{1}{4}$

So $\frac{1}{x^2-4} = \frac{1}{4(x-2)} - \frac{1}{4(x+2)}$

$\int_3^\infty \frac{1}{x^2-4}\,dx = \lim_{k\to\infty} \int_3^k \frac{1}{4}\left(\frac{1}{x-2} - \frac{1}{x+2}\right) dx$

$= \lim_{k\to\infty} \frac{1}{4}[\ln(x-2) - \ln(x+2)]_3^k \qquad (x-2 > 0 \text{ for } x \geqslant 3)$

$= \lim_{k\to\infty} \frac{1}{4}[\ln(\frac{x-2}{x+2})]_3^k$

$= \lim_{k\to\infty} \frac{1}{4}\left(\ln\left(\frac{k-2}{k+2}\right) - \ln\left(\frac{1}{5}\right)\right)$

As $k \to \infty$, $\frac{k-2}{k+2} = \frac{1-2/k}{1+2/k} \to 1$,

so $\ln\left(\frac{k-2}{k+2}\right) \to \ln 1 = 0$

So $\int_3^\infty \frac{1}{x^2-4}\,dx = \frac{1}{4}\left(0 - \ln\frac{1}{5}\right) = \frac{1}{4}\ln 5$

$m = 4, \quad n = 5$

© OCR 2025

stemup.co.uk

8(a)

When $n = 1$: $f'(x) = \frac{1}{1+x}$

Formula with $n = 1$: $(-1)^2(0)!(1+x)^{-1} = \frac{1}{1+x}$

These agree, so the result is true for $n = 1$.

Assume the result is true for $n = k$:

$f^{(k)}(x) = (-1)^{k+1}(k-1)! \,/\, (1+x)^k = (-1)^{k+1}(k-1)!(1+x)^{-k}$

Then $f^{(k+1)}(x) = \frac{d}{dx}[(-1)^{k+1}(k-1)!(1+x)^{-k}]$

$= (-1)^{k+1}(k-1)! \times (-k)(1+x)^{-k-1}$

$= (-1)^{k+2} \times k(k-1)! \times (1+x)^{-(k+1)}$

$= (-1)^{(k+1)+1}((k+1)-1)! \,/\, (1+x)^{k+1}$

which is the result with $n = k + 1$.

So if the result is true for $n = k$, it is true for $n = k + 1$.

Since it is true for $n = 1$, by mathematical induction it is true for all integers $n \geq 1$.

© OCR 2025

8(b)	$f(0) = \ln 1 = 0$
	From (a): $f^{(n)}(0) = (-1)^{n+1}(n-1)! \,/\, 1^n = (-1)^{n+1}(n-1)!$
	e.g. $f'(0) = 1$, $f''(0) = -1$, $f'''(0) = 2!$
	Maclaurin: $f(x) = f(0) + f'(0)x + \frac{f''(0)}{2!}x^2 + \frac{f'''(0)}{3!}x^3 + \ldots$
	Coefficient of x^n: $f^{(n)}(0) \,/\, n! = (-1)^{n+1}(n-1)! \,/\, n! = (-1)^{n+1} \,/\, n$
	$\ln(1+x) = 0 + x - \frac{1!}{2!}x^2 + \frac{2!}{3!}x^3 - \ldots + ((-1)^{n+1}/n)x^n + \ldots$
	$\ln(1+x) = x - \frac{x^2}{2} + \frac{x^3}{3} - \ldots + \frac{(-1)^{n+1}x^n}{n} + \ldots$
	for $-1 < x \leqslant 1$, as required

DO NOT WRITE IN THIS SPACE

© OCR 2025

stemup.co.uk

9(a)(i)

C: w^2, D: w^3, E: w^4

9(a)(ii)

$z^5 = 1$

9(a)(iii)

Sum of roots $= 1 + w + w^2 + w^3 + w^4$ (geometric series, ratio $w \neq 1$)

$= \frac{1 - w^5}{1 - w}$

$= 0$, since $w^5 = 1$

9(b)(i)

$w = \cos(\frac{2\pi}{5}) + i\sin(\frac{2\pi}{5})$ $(r = 1,\ k = \frac{2}{5})$

9(b)(ii)

$AB = |w - 1|$

$w - 1 = (\cos\frac{2\pi}{5} - 1) + i\sin\frac{2\pi}{5}$

$|w - 1|^2 = (\cos\frac{2\pi}{5} - 1)^2 + \sin^2\frac{2\pi}{5}$

$= \cos^2\frac{2\pi}{5} - 2\cos\frac{2\pi}{5} + 1 + \sin^2\frac{2\pi}{5}$

$= 2 - 2\cos\frac{2\pi}{5}$

Using $\cos 2A = 1 - 2\sin^2 A$ with $A = \frac{\pi}{5}$:

$= 2 - 2(1 - 2\sin^2\frac{\pi}{5}) = 4\sin^2\frac{\pi}{5}$

$AB = 2\sin\frac{\pi}{5}$ (positive since $0 < \frac{\pi}{5} < \frac{\pi}{2}$)

The pentagon is regular, so each side has length $2\sin\frac{\pi}{5}$, as required

© OCR 2025

10

$x^2 - x + 1 = (x - \frac{1}{2})^2 + \frac{3}{4}$

$\int_0^{1/2} \frac{2}{x^2 - x + 1}\, dx$

$= 2\int_0^{1/2} \frac{1}{(x - 1/2)^2 + 3/4}\, dx$

Using $\int \frac{1}{u^2 + a^2}\, du = \frac{1}{a} \arctan(\frac{u}{a})$ with $a = \frac{\sqrt{3}}{2}$:

$= 2 \times \frac{2}{\sqrt{3}} [\arctan(\frac{x - 1/2}{\sqrt{3}/2})]_0^{1/2}$

$= \frac{4}{\sqrt{3}} [\arctan(\frac{2x - 1}{\sqrt{3}})]_0^{1/2}$

$= \frac{4}{\sqrt{3}} (\arctan 0 - \arctan(-\frac{1}{\sqrt{3}}))$

$= \frac{4}{\sqrt{3}} (0 - (-\frac{\pi}{6}))$

$= \frac{4\pi}{6\sqrt{3}}$

$= \frac{2\pi}{3\sqrt{3}} \quad (= \frac{2\sqrt{3}}{9}\pi)$

© OCR 2025

stemup.co.uk

11(a)

$c = 0$: l_2: $\mathbf{r} = 2\mathbf{i} + \mathbf{j} + 3\mathbf{k} + \lambda(-\mathbf{i} + 2\mathbf{k})$

l_1 passes through $(3, -1, 2)$ with direction $a\mathbf{i} + b\mathbf{j} + \mathbf{k}$

Perpendicular: $(a\mathbf{i} + b\mathbf{j} + \mathbf{k})\cdot(-\mathbf{i} + 0\mathbf{j} + 2\mathbf{k}) = 0$

$-a + 2 = 0 \Rightarrow a = 2$

l_1: $x = 3 + 2\mu$, $y = -1 + b\mu$, $z = 2 + \mu$

l_2: $x = 2 - \lambda$, $y = 1$, $z = 3 + 2\lambda$

x: $3 + 2\mu = 2 - \lambda \Rightarrow \lambda + 2\mu = -1$

z: $2 + \mu = 3 + 2\lambda \Rightarrow 2\lambda - \mu = -1 \Rightarrow \mu = 2\lambda + 1$

$\lambda + 2(2\lambda + 1) = -1 \Rightarrow 5\lambda = -3 \Rightarrow \lambda = -\frac{3}{5}$

$\mu = 2(-\frac{3}{5}) + 1 = -\frac{1}{5}$

(y: $-1 + b(-\frac{1}{5}) = 1$ gives $b = -10$, so the lines do meet)

Using l_2 with $\lambda = -\frac{3}{5}$:

$x = 2 + \frac{3}{5} = \frac{13}{5}$, $y = 1$, $z = 3 - \frac{6}{5} = \frac{9}{5}$

Point of intersection: $(\frac{13}{5}, 1, \frac{9}{5})$

© OCR 2025

11(b)(i)

$c = 4$: direction of l_2 is $-i + 4j + 2k$

Parallel: $ai + bj + k = t(-i + 4j + 2k)$

k: $1 = 2t \Rightarrow t = \frac{1}{2}$

$a = -\frac{1}{2}$, $b = 4 \times \frac{1}{2} = 2$

11(b)(ii)

Point $A(3, -1, 2)$ on l_1, point $B(2, 1, 3)$ on l_2

$\overrightarrow{AB} = (2 - 3)i + (1 + 1)j + (3 - 2)k = -i + 2j + k$

Direction $d = -i + 4j + 2k$

$\overrightarrow{AB} \times d = (2 \times 2 - 1 \times 4)i - ((-1)(2) - (1)(-1))j + ((-1)(4) - (2)(-1))k$

$= 0i - (-1)j + (-2)k = j - 2k$

$|\overrightarrow{AB} \times d| = \sqrt{0^2 + 1^2 + (-2)^2} = \sqrt{5}$

$|d| = \sqrt{(-1)^2 + 4^2 + 2^2} = \sqrt{21}$

Distance $= |\overrightarrow{AB} \times d| \div |d|$

$= \frac{\sqrt{5}}{\sqrt{21}} = \sqrt{(5/21)}$

$= 0.488$ (3 s.f.)

© OCR 2025

stemup.co.uk

12

Sum of roots: $\alpha + \frac{2}{\alpha} + \beta + (-\beta) = -\frac{(-1)}{1} = 1$

$\alpha + \frac{2}{\alpha} = 1 \Rightarrow \alpha^2 - \alpha + 2 = 0$

$\alpha = \frac{1 \pm \sqrt{(1-8)}}{2} = \frac{1 \pm i\sqrt{7}}{2}$

α and $\frac{2}{\alpha}$ are the two roots of $\alpha^2 - \alpha + 2 = 0$ (product 2),

so they are $\frac{1}{2} + \frac{\sqrt{7}}{2}i$ and $\frac{1}{2} - \frac{\sqrt{7}}{2}i$

Product of roots: $\alpha \times \frac{2}{\alpha} \times \beta \times (-\beta) = \frac{18}{1} = 18$

$-2\beta^2 = 18 \Rightarrow \beta^2 = -9 \Rightarrow \beta = 3i, \quad -\beta = -3i$

Roots $\frac{1}{2} \pm \frac{\sqrt{7}}{2}i$ give the factor $z^2 - z + 2$

Roots $\pm 3i$ give the factor $z^2 + 9$

$(z^2 - z + 2)(z^2 + 9) = z^4 - z^3 + 2z^2 + 9z^2 - 9z + 18$

$= z^4 - z^3 + 11z^2 - 9z + 18$

Comparing with $z^4 - z^3 + cz^2 + dz + 18$: $c = 11, \quad d = -9$

The four roots of the equation: $\frac{1}{2} + \frac{\sqrt{7}}{2}i, \quad \frac{1}{2} - \frac{\sqrt{7}}{2}i,$

$3i, \quad -3i$

$c =$ 11

$d =$ −9

© OCR 2025

13(a)

Divide by x: $\frac{dy}{dx} - \frac{2}{x}y = \frac{2 + x^2}{x}$

Integrating factor $= e^{\int -2/x\, dx}$

$= e^{-2 \ln x} = e^{\ln(x^{-2})}$

$= x^{-2}$, as required

13(b)

Multiply by x^{-2}: $x^{-2}\frac{dy}{dx} - 2x^{-3}y = \frac{2 + x^2}{x^3}$

$\frac{d}{dx}(yx^{-2}) = 2x^{-3} + x^{-1}$

$yx^{-2} = \int (2x^{-3} + x^{-1})\, dx = -x^{-2} + \ln x + c$

$y = -1 + x^2 \ln x + cx^2$

Through $(1, 0)$: $0 = -1 + 0 + c \quad \Rightarrow \quad c = 1$

$y = x^2 \ln x + x^2 - 1$

$\frac{dy}{dx} = 2x \ln x + x^2 \times \frac{1}{x} + 2x = 2x \ln x + 3x$

Stationary point: $\frac{dy}{dx} = 0 \quad \Rightarrow \quad x(2 \ln x + 3) = 0$

$x > 0$ ($x = 0$ is not in the domain of $\ln x$), so $2 \ln x + 3 = 0$

$\ln x = -\frac{3}{2}$

$x = e^{-3/2}$

© OCR 2025

stemup.co.uk

14(a)

$\cosh^2 x + \sinh^2 x = \frac{1}{4}(e^x + e^{-x})^2 + \frac{1}{4}(e^x - e^{-x})^2$

$= \frac{1}{4}(e^{2x} + 2 + e^{-2x}) + \frac{1}{4}(e^{2x} - 2 + e^{-2x})$

$= \frac{1}{4}(2e^{2x} + 2e^{-2x})$

$= \frac{1}{2}(e^{2x} + e^{-2x}) = \cosh 2x$, as required

14(b)

$M\begin{pmatrix}0\\0\end{pmatrix} = \begin{pmatrix}0\\0\end{pmatrix}$ $M\begin{pmatrix}1\\0\end{pmatrix} = \begin{pmatrix}\cosh x\\ \sinh x\end{pmatrix}$

$M\begin{pmatrix}0\\1\end{pmatrix} = \begin{pmatrix}\sinh x\\ \cosh x\end{pmatrix}$ $M\begin{pmatrix}1\\1\end{pmatrix} = \begin{pmatrix}\cosh x + \sinh x\\ \sinh x + \cosh x\end{pmatrix}$

Images: $O(0, 0)$, $P(\cosh x, \sinh x)$, $Q(\sinh x, \cosh x)$,

$R(\cosh x + \sinh x, \cosh x + \sinh x)$

$OP^2 = \cosh^2 x + \sinh^2 x$

$OQ^2 = \sinh^2 x + \cosh^2 x$

$\overrightarrow{PR} = (\sinh x, \cosh x)$ so $PR^2 = \sinh^2 x + \cosh^2 x$

$\overrightarrow{QR} = (\cosh x, \sinh x)$ so $QR^2 = \cosh^2 x + \sinh^2 x$

So $OP = PR = RQ = QO$: all four sides are equal,

so the image $OPRQ$ is a rhombus.

$\det M = \cosh^2 x - \sinh^2 x = 1$

The area scale factor of T is $|\det M| = 1$, so T preserves area.

The unit square has area 1, so the rhombus has area $1 \times 1 = 1$,

so the rhombus has unit area, as required.

(answer space continued on next page)

© OCR 2025

14(b) **(continued)**

14(c)

Side length: $OP^2 = \cosh^2 x + \sinh^2 x = \cosh 2x$ (from (a))

$\cosh 2x = 2^2 = 4$

$2x = \operatorname{arcosh} 4 = \ln(4 + \sqrt{4^2 - 1})$ $(x > 0)$

$2x = \ln(4 + \sqrt{15})$

$x = \frac{1}{2} \ln(4 + \sqrt{15})$

© OCR 2025

stemup.co.uk

15(a)

$(3 - e^{4i\theta})(3 - e^{-4i\theta}) = 9 - 3e^{-4i\theta} - 3e^{4i\theta} + e^{0}$

$= 10 - 3(e^{4i\theta} + e^{-4i\theta}) = 10 - 3(2\cos 4\theta)$

$= 10 - 6\cos 4\theta \qquad (a = 10, \ b = -6)$

15(b)

$C + iS = (\cos\theta + i\sin\theta) + \frac{1}{3}(\cos 5\theta + i\sin 5\theta) + \frac{1}{9}(\cos 9\theta + i\sin 9\theta) + \ldots$

$= e^{i\theta} + \frac{1}{3}e^{5i\theta} + \frac{1}{9}e^{9i\theta} + \frac{1}{27}e^{13i\theta} + \ldots$

$= e^{i\theta} + \frac{1}{3}e^{i\theta}(e^{i\theta})^{4} + \frac{1}{9}e^{i\theta}(e^{i\theta})^{8} + \ldots$

Geometric series: first term $e^{i\theta}$, common ratio $\frac{1}{3}e^{4i\theta}$

$\left|\frac{1}{3}e^{4i\theta}\right| = \frac{1}{3} < 1$, so the sum to infinity exists

$C + iS = \frac{e^{i\theta}}{1 - (1/3)e^{4i\theta}}$

$= \frac{3e^{i\theta}}{3 - e^{4i\theta}}$, as required

15(c)

$C + iS = \frac{3e^{i\theta}(3 - e^{-4i\theta})}{(3 - e^{4i\theta})(3 - e^{-4i\theta})}$

$= \frac{9e^{i\theta} - 3e^{-3i\theta}}{10 - 6\cos 4\theta}$ (using (a))

$= \frac{9(\cos\theta + i\sin\theta) - 3(\cos 3\theta - i\sin 3\theta)}{10 - 6\cos 4\theta}$

$C = \mathrm{Re}(C + iS)$

$C = \frac{9\cos\theta - 3\cos 3\theta}{10 - 6\cos 4\theta}$, as required

© OCR 2025

16(a)

$A = \int_0^4 \frac{x+3}{\sqrt{(x^2+9)}}\,dx$

$= \int_0^4 \frac{x}{\sqrt{(x^2+9)}}\,dx + \int_0^4 \frac{3}{\sqrt{(x^2+9)}}\,dx$

$\int x(x^2+9)^{-1/2}\,dx = (x^2+9)^{1/2} = \sqrt{x^2+9}$

$\int \frac{3}{\sqrt{(x^2+9)}}\,dx = 3\,\text{arsinh}(\frac{x}{3}) = 3\ln(x+\sqrt{x^2+9}) + \text{const}$

$A = [\sqrt{x^2+9} + 3\ln(x+\sqrt{x^2+9})]_0^4$

$= (\sqrt{25} + 3\ln(4+5)) - (\sqrt{9} + 3\ln(0+3))$

$= 5 + 3\ln 9 - 3 - 3\ln 3$

$= 2 + 3\ln 3$

$= 2 + \ln 27$

$p = 2, \quad q = 27$

© OCR 2025

stemup.co.uk

16(b)

$V = \pi\int_0^4 y^2\,dx = \pi\int_0^4 \frac{(x+3)^2}{x^2+9}\,dx$

$(x+3)^2 = x^2 + 6x + 9$, so $\frac{(x+3)^2}{x^2+9} = 1 + \frac{6x}{x^2+9}$

$V = \pi\int_0^4 \left(1 + \frac{6x}{x^2+9}\right)dx$

$= \pi[x + 3\ln(x^2+9)]_0^4$

$= \pi[(4 + 3\ln 25) - (0 + 3\ln 9)]$

$= \pi\left(4 + 3\ln\left(\frac{25}{9}\right)\right)$

$a = 4, \quad b = 3, \quad c = 25, \quad d = 9$

© OCR 2025

17(a)(i)

$\frac{dh}{dt} = 0.2(20 - h)$

17(a)(ii)

Growth stops when $\frac{dh}{dt} = 0$: $0.2(20 - h) = 0 \Rightarrow h = 20$

(for $h < 20$, $\frac{dh}{dt} > 0$, so h increases towards 20) so max height is 20 m

17(a)(iii)

$\int \frac{1}{20 - h}\, dh = \int 0.2\, dt$

$-\ln(20 - h) = 0.2t + c \qquad (20 - h > 0)$

$\ln(20 - h) = -0.2t - c$

$20 - h = Ae^{-0.2t}$

When $t = 0$, $h = 0$: $20 - 0 = A \Rightarrow A = 20$

$h = 20 - 20e^{-0.2t}$

$h = 20(1 - e^{-0.2t})$, as required

17(a)(iv)

$t = 5$: $h = 20(1 - e^{-1}) = 12.64$ m (4 s.f.)

Maximum height would be $2 \times 12.64 = 25.28$ m $\neq 20$ m

(equivalently $12.64 \neq 10$, half of 20)

So this model is not consistent with a maximum height of 20 m.

© OCR 2025

stemup.co.uk

17(b)(i)

Auxiliary equation: $\lambda^2 + 0.3\lambda + 0.02 = 0$

$(\lambda + 0.1)(\lambda + 0.2) = 0 \Rightarrow \lambda = -0.1, -0.2$

Complementary function: $h = Ae^{-0.1t} + Be^{-0.2t}$

Particular integral: try $h = k$: $0 + 0 + 0.02k = 0.4 \Rightarrow k = 20$

General solution: $h = Ae^{-0.1t} + Be^{-0.2t} + 20$

17(b)(ii)

As $t \to \infty$, $e^{-0.1t} \to 0$ and $e^{-0.2t} \to 0$, so $h \to 20$

So the refined model also predicts a maximum height of 20 m.

17(b)(iii)

$t = 0, h = 0$: $A + B + 20 = 0 \Rightarrow A + B = -20$

$\frac{dh}{dt} = -0.1Ae^{-0.1t} - 0.2Be^{-0.2t}$

$t = 0, \frac{dh}{dt} = 2.9$: $-0.1A - 0.2B = 2.9 \Rightarrow A + 2B = -29$

Subtracting: $B = -9$, so $A = -20 - (-9) = -11$

Particular solution: $h = 20 - 11e^{-0.1t} - 9e^{-0.2t}$

$t = 5$: $h = 20 - 11e^{-0.5} - 9e^{-1}$

$= 20 - 6.6718... - 3.3109...$

$= 10.017...$ m

Twice this is $2 \times 10.017 = 20.03 \approx 20$ m ($h(5) \approx 10$, half of 20)

So the refined model is consistent with a maximum height of 20 m.

© OCR 2025

EXTRA ANSWER SPACE

If you need extra space use these lined pages. You must write the question numbers clearly in the margin.

© OCR 2025

stemup.co.uk

OCR
Oxford Cambridge and RSA

Copyright Information

OCR is committed to seeking permission to reproduce all third-party content that it uses in its assessment materials. OCR has attempted to identify and contact all copyright holders whose work is used in this paper. To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced in the OCR Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download from our public website (www.ocr.org.uk) after the live examination series.

If OCR has unwittingly failed to correctly acknowledge or clear any third-party content in this assessment material, OCR will be happy to correct its mistake at the earliest possible opportunity.

For queries or further information please contact The OCR Copyright Team, The Triangle Building, Shaftesbury Road, Cambridge CB2 8EA.

OCR is part of Cambridge University Press & Assessment, which is itself a department of the University of Cambridge.

© OCR 2025

stemup.co.uk