OCR
Oxford Cambridge and RSA

Thursday 20 June 2024 – Afternoon

A Level Mathematics B (MEI)

H640/03 Pure Mathematics and Comprehension

Time allowed: 2 hours

136602435 1

You must have:
- the Printed Answer Booklet
- the Insert
- a scientific or graphical calculator

QP

INSTRUCTIONS
- Use black ink. You can use an HB pencil, but only for graphs and diagrams.
- Write your answer to each question in the space provided in the **Printed Answer Booklet**. If you need extra space use the lined pages at the end of the Printed Answer Booklet. The question numbers must be clearly shown.
- Fill in the boxes on the front of the Printed Answer Booklet.
- Answer **all** the questions.
- Where appropriate, your answer should be supported with working. Marks might be given for using a correct method, even if your answer is wrong.
- Give your final answers to a degree of accuracy that is appropriate to the context.
- Do **not** send this Question Paper for marking. Keep it in the centre or recycle it.

INFORMATION
- The total mark for this paper is **75**.
- The marks for each question are shown in brackets **[ ]**.
- This document has **12** pages.

ADVICE
- Read each question carefully before you start your answer.

© OCR 2024 [603/1002/9]
DC (DE/SG) 330047/5
OCR is an exempt Charity

stemup.co.uk

Formulae A Level Mathematics B (MEI) (H640)

Arithmetic series

$S_n = \frac{1}{2}n(a+l) = \frac{1}{2}n\{2a+(n-1)d\}$

Geometric series

$S_n = \dfrac{a(1-r^n)}{1-r}$

$S_\infty = \dfrac{a}{1-r}$ for $|r| < 1$

Binomial series

$(a+b)^n = a^n + {}^n\mathrm{C}_1 a^{n-1}b + {}^n\mathrm{C}_2 a^{n-2}b^2 + \ldots + {}^n\mathrm{C}_r a^{n-r}b^r + \ldots + b^n \qquad (n \in \mathbb{N}),$

where ${}^n\mathrm{C}_r = {}_n\mathrm{C}_r = \binom{n}{r} = \dfrac{n!}{r!(n-r)!}$

$(1+x)^n = 1 + nx + \dfrac{n(n-1)}{2!}x^2 + \ldots + \dfrac{n(n-1)\ldots(n-r+1)}{r!}x^r + \ldots \qquad (|x| < 1,\ n \in \mathbb{R})$

Differentiation

$\mathrm{f}(x)$	$\mathrm{f}'(x)$
$\tan kx$	$k\sec^2 kx$
$\sec x$	$\sec x \tan x$
$\cot x$	$-\mathrm{cosec}^2 x$
$\mathrm{cosec}\, x$	$-\mathrm{cosec}\, x \cot x$

Quotient Rule $y = \dfrac{u}{v},\ \dfrac{\mathrm{d}y}{\mathrm{d}x} = \dfrac{v\frac{\mathrm{d}u}{\mathrm{d}x} - u\frac{\mathrm{d}v}{\mathrm{d}x}}{v^2}$

Differentiation from first principles

$\mathrm{f}'(x) = \lim_{h\to 0}\dfrac{\mathrm{f}(x+h)-\mathrm{f}(x)}{h}$

Integration

$\displaystyle\int \frac{\mathrm{f}'(x)}{\mathrm{f}(x)}\,\mathrm{d}x = \ln|\mathrm{f}(x)| + c$

$\displaystyle\int \mathrm{f}'(x)\big(\mathrm{f}(x)\big)^n\,\mathrm{d}x = \frac{1}{n+1}\big(\mathrm{f}(x)\big)^{n+1} + c$

Integration by parts $\displaystyle\int u\frac{\mathrm{d}v}{\mathrm{d}x}\,\mathrm{d}x = uv - \int v\frac{\mathrm{d}u}{\mathrm{d}x}\,\mathrm{d}x$

Small angle approximations

$\sin\theta \approx \theta,\ \cos\theta \approx 1 - \frac{1}{2}\theta^2,\ \tan\theta \approx \theta$ where θ is measured in radians

© OCR 2024

stemup.co.uk

Trigonometric identities

$\sin(A \pm B) = \sin A \cos B \pm \cos A \sin B$

$\cos(A \pm B) = \cos A \cos B \mp \sin A \sin B$

$\tan(A \pm B) = \dfrac{\tan A \pm \tan B}{1 \mp \tan A \tan B} \qquad \left(A \pm B \neq (k + \frac{1}{2})\pi\right)$

Numerical methods

Trapezium rule: $\displaystyle\int_a^b y\,\mathrm{d}x \approx \tfrac{1}{2}h\{(y_0 + y_n) + 2(y_1 + y_2 + \ldots + y_{n-1})\}$, where $h = \dfrac{b-a}{n}$

The Newton-Raphson iteration for solving $\mathrm{f}(x) = 0$: $x_{n+1} = x_n - \dfrac{\mathrm{f}(x_n)}{\mathrm{f}'(x_n)}$

Probability

$\mathrm{P}(A \cup B) = \mathrm{P}(A) + \mathrm{P}(B) - \mathrm{P}(A \cap B)$

$\mathrm{P}(A \cap B) = \mathrm{P}(A)\mathrm{P}(B \mid A) = \mathrm{P}(B)\mathrm{P}(A \mid B)$ **or** $\mathrm{P}(A \mid B) = \dfrac{\mathrm{P}(A \cap B)}{\mathrm{P}(B)}$

Sample variance

$s^2 = \dfrac{1}{n-1}S_{xx}$ where $S_{xx} = \Sigma(x_i - \bar{x})^2 = \Sigma x_i^2 - \dfrac{(\Sigma x_i)^2}{n} = \Sigma x_i^2 - n\bar{x}^2$

Standard deviation, $s = \sqrt{\text{variance}}$

The binomial distribution

If $X \sim \mathrm{B}(n, p)$ then $\mathrm{P}(X = r) = {}^n\mathrm{C}_r p^r q^{n-r}$ where $q = 1 - p$


Mean of X is np

Hypothesis testing for the mean of a Normal distribution

If $X \sim \mathrm{N}(\mu, \sigma^2)$ then $\overline{X} \sim \mathrm{N}\left(\mu, \dfrac{\sigma^2}{n}\right)$ and $\dfrac{\overline{X} - \mu}{\sigma/\sqrt{n}} \sim \mathrm{N}(0, 1)$

Percentage points of the Normal distribution

p	10	5	2	1
z	1.645	1.960	2.326	2.576



Kinematics

Motion in a straight line	Motion in two dimensions
$v = u + at$	$\mathbf{v} = \mathbf{u} + \mathbf{a}t$
$s = ut + \frac{1}{2}at^2$	$\mathbf{s} = \mathbf{u}t + \frac{1}{2}\mathbf{a}t^2$
$s = \frac{1}{2}(u + v)t$	$\mathbf{s} = \frac{1}{2}(\mathbf{u} + \mathbf{v})t$
$v^2 = u^2 + 2as$	
$s = vt - \frac{1}{2}at^2$	$\mathbf{s} = \mathbf{v}t - \frac{1}{2}\mathbf{a}t^2$

Section A (60 marks)

1 Solve the inequality $\frac{x}{5} > 6 - x$. **[2]**

2 **(a)** The function $f(x)$ is defined by

$f(x) = \sqrt{1+2x}$ for $x \geqslant -\frac{1}{2}$.


Find an expression for $f^{-1}(x)$ and state the domain of this inverse function. **[3]**

(b) Explain why $g(x) = 1 + x^2$, with domain all real numbers, has no inverse function. **[1]**

3 **In this question you must show detailed reasoning.**

The diagram shows the curve with equation $y = x^5$ and the square OABC where the points A, B and C have coordinates (1, 0), (1, 1) and (0, 1) respectively.

The curve cuts the square into two parts.



Show that the relationship between the areas of the two parts of the square is

$\frac{\text{Area to left of curve}}{\text{Area below curve}} = 5$. **[4]**

4 **In this question you must show detailed reasoning.**

Determine the exact value of $\frac{1}{\sqrt{2}+1} + \frac{1}{\sqrt{3}+\sqrt{2}} + \frac{1}{2+\sqrt{3}}$. **[2]**

stemup.co.uk

5 **In this question you must show detailed reasoning.**

Using the substitution $u = x + 1$, find the value of the positive integer c such that

$$\int_{c}^{c+4} \frac{x}{(x+1)^2}\,\mathrm{d}x = \ln 3 - \frac{1}{3}.$$ **[6]**

6 **In this question you must show detailed reasoning.**

Solve the equation $\tan x - 3\cot x = 2$ for values of x in the interval $0° \leqslant x \leqslant 360°$. **[5]**

7 Prove that $\sin 8\theta \tan 4\theta + \cos 8\theta = 1$. **[3]**

8 **In this question you must show detailed reasoning.**

(a) Express $\cos x + \sqrt{3}\sin x$ in the form $R\sin(x+\alpha)$, where $R > 0$ and $0 < \alpha < \frac{1}{2}\pi$. Give the values of R and α in exact form. **[4]**

(b) Hence solve the equation $\cos x = \sqrt{3}(1 - \sin x)$ for values of x in the interval $-\pi \leqslant x \leqslant \pi$. Give the roots of this equation in exact form. **[4]**


© OCR 2024

stemup.co.uk

9 This question is about the equation $f(x) = 0$, where $f(x) = x^4 - x - \frac{1}{3x-2}$.

Fig. 9.1 shows the curve $y = f(x)$.

Fig. 9.1



(a) Show, by calculation, that the equation $f(x) = 0$ has a root between $x = 1$ and $x = 2$. **[2]**

(b) **Fig. 9.2** shows part of a spreadsheet being used to find a root of the equation.

Fig. 9.2

	A	B
1	x	f(x)
2	1.5	3.1625
3	1.25	0.619977679
4	1.125	-0.250466087
5		

Write down a suitable number to use as the next value of x in the spreadsheet. **[1]**

(c) Determine a root of the equation $f(x) = 0$. Give your answer correct to **1** decimal place. **[1]**

© OCR 2024

stemup.co.uk

(d) **Fig. 9.3** shows a similar spreadsheet being used to search for another root of $f(x) = 0$.


Fig. 9.3

	A	B
1	x	f(x)
2	0	0.5
3	1	-1
4	0.5	1.5625
5	0.75	-4.4336
6	0.6	4.5296
7	0.7	-10.4599
8	0.65	19.5285
9	0.675	-40.4674
10	0.6625	79.5301
11	0.66875	-160.4687

(i) Explain why it looks from rows 2 and 3 of the spreadsheet as if there is a root between 0 and 1. **[1]**

(ii) Explain why this process will **not** find a root between 0 and 1. **[1]**

10 The diagram below shows the curve $y = f(x)$.




Sketch the graph of the gradient function, $y = f'(x)$, on the copy of the diagram in the **Printed Answer Booklet**. **[3]**

stemup.co.uk

11 **Fig. 11.1** shows the curve with equation $y = g(x)$ where $g(x) = x\sin x + \cos x$ and the curve of the gradient function $y = g'(x)$ for $-2\pi \leqslant x \leqslant 2\pi$.

Fig. 11.1




(a) Show that the x-coordinates of the points on the curve $y = g(x)$ where the gradient is 1 satisfy the equation $\frac{1}{x} - \cos x = 0$. **[3]**

stemup.co.uk

Fig. 11.2 shows part of the curve with equation $y = \frac{1}{x} - \cos x$.

Fig. 11.2



(b) Use the Newton-Raphson method with a suitable starting value to find the smallest positive x-coordinate of a point on the curve $y = x\sin x + \cos x$ where the gradient is 1.

You should write down at least the following.


- The iteration you use
- The starting value
- The solution correct to **4** decimal places **[4]**

(c) Explain why $x_1 = 3$ is **not** a suitable starting value for the Newton-Raphson method in part **(b)**. **[1]**

stemup.co.uk

12 The diagram shows the curve with parametric equations

$x = \sin 2\theta + 2,\ y = 2\cos\theta + \cos 2\theta,$ for $0 \leqslant \theta < 2\pi$.



(a) In this question you must show detailed reasoning.

Determine the exact coordinates of all the stationary points on the curve. **[8]**

(b) Write down the equation of the line of symmetry of the curve. **[1]**

© OCR 2024

stemup.co.uk

Section B (15 marks)

The questions in this section refer to the article on the Insert. You should read the article before attempting the questions.

13 Substitute appropriate values of t_1 and t_2 to verify that $t_1 t_2$ gives the correct value for the y-coordinate of the point of intersection of the tangents at the points A and B in **Fig. C1**. **[1]**

14 Substitute appropriate values of t_1 and t_2 to verify that the expression $t_1^2 + t_2^2 + t_1 t_2 + \frac{1}{2}$ gives the correct value for the y-coordinate of the point of intersection of the normals at the points A and B in **Fig. C2**. **[1]**

15 **(a)** Show that, for the curve $y = ax^2 + bx + c$, the equation of the tangent at the point with x-coordinate t is $y = (2at + b)x - at^2 + c$. **[3]**

(b) Hence show that for the curve with equation $y = ax^2 + bx + c$, the tangents at two points, P and Q, on the curve cross at a point which has x-coordinate equal to the mean of the x-coordinates of points P and Q, as given in lines 11 to 14. **[3]**

16 Show that the expression $a\left(\frac{x_P + x_Q}{2}\right)^2 + b\left(\frac{x_P + x_Q}{2}\right) + c - a\left(\frac{x_P - x_Q}{2}\right)^2$ is equivalent to $ax_P x_Q + b\left(\frac{x_P + x_Q}{2}\right) + c$, as given in lines 15 and 16. **[2]**

17 Show that, for the curve $y = x^2$, the equation of the normal at the point (t, t^2) is $y = -\frac{x}{2t} + t^2 + \frac{1}{2}$, as given in line 27. **[3]**

18 A student is investigating the intersection points of tangents to the curve $y = 6x^2 - 7x + 1$. She uses software to draw tangents at pairs of points with x-coordinates differing by 5.

Find the equation of the curve that all the intersection points lie on. **[2]**

END OF QUESTION PAPER

© OCR 2024

stemup.co.uk

OCR
Oxford Cambridge and RSA

Copyright Information

OCR is committed to seeking permission to reproduce all third-party content that it uses in its assessment materials. OCR has attempted to identify and contact all copyright holders whose work is used in this paper. To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced in the OCR Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download from our public website (www.ocr.org.uk) after the live examination series.

If OCR has unwittingly failed to correctly acknowledge or clear any third-party content in this assessment material, OCR will be happy to correct its mistake at the earliest possible opportunity.

For queries or further information please contact The OCR Copyright Team, The Triangle Building, Shaftesbury Road, Cambridge CB2 8EA.

OCR is part of Cambridge University Press & Assessment, which is itself a department of the University of Cambridge.

© OCR 2024

stemup.co.uk

Surname	StemUp Model Solution
Answer booklet	Write the question number in the left-hand margin at the start of each answer.

Question number

1

$\frac{x}{5} > 6 - x \quad \Rightarrow \quad x > 30 - 5x$

$6x > 30$

$x > 5$

2 (a)

$y = \sqrt{1 + 2x} \quad \Rightarrow \quad y^2 = 1 + 2x \quad \Rightarrow \quad x = \frac{y^2 - 1}{2}$

$f^{-1}(x) = \frac{x^2 - 1}{2}$

Domain of f^{-1}: $x \geq 0$ (the range of f is $f(x) \geq 0$)

2 (b)

g is many-to-one, e.g. $g(2) = g(-2) = 5$, so it is not one-to-one and therefore has no inverse function.

3

Area below curve $= \int_0^1 x^5 \, dx = \left[\frac{x^6}{6} \right]_0^1 = \frac{1}{6} - 0 = \frac{1}{6}$

Area to left of curve = area of square $- \frac{1}{6} = 1 - \frac{1}{6} = \frac{5}{6}$

$\frac{\text{Area to left of curve}}{\text{Area below curve}} = \frac{5}{6} \div \frac{1}{6} = \frac{5}{6} \times 6 = 5$ as required

4

$\frac{1}{\sqrt{2} + 1} = \frac{\sqrt{2} - 1}{(\sqrt{2} + 1)(\sqrt{2} - 1)} = \frac{\sqrt{2} - 1}{2 - 1} = \sqrt{2} - 1$

$\frac{1}{\sqrt{3} + \sqrt{2}} = \frac{\sqrt{3} - \sqrt{2}}{(\sqrt{3} + \sqrt{2})(\sqrt{3} - \sqrt{2})} = \frac{\sqrt{3} - \sqrt{2}}{3 - 2} = \sqrt{3} - \sqrt{2}$

$\frac{1}{2 + \sqrt{3}} = \frac{2 - \sqrt{3}}{(2 + \sqrt{3})(2 - \sqrt{3})} = \frac{2 - \sqrt{3}}{4 - 3} = 2 - \sqrt{3}$

Sum $= (\sqrt{2} - 1) + (\sqrt{3} - \sqrt{2}) + (2 - \sqrt{3}) = 2 - 1 = 1$

5

$u = x + 1 \quad \Rightarrow \quad \frac{du}{dx} = 1$, so $du = dx$ and $x = u - 1$

$x = c \quad \Rightarrow \quad u = c + 1; \qquad x = c + 4 \quad \Rightarrow \quad u = c + 5$

Question number

$$\int_c^{c+4} \frac{x}{(x+1)^2}\,dx = \int_{c+1}^{c+5} \frac{u-1}{u^2}\,du = \int_{c+1}^{c+5} \left(\frac{1}{u} - \frac{1}{u^2}\right) du$$

$$= \left[\ln u + \frac{1}{u}\right]_{c+1}^{c+5}$$

$$= \ln(c+5) + \frac{1}{c+5} - \ln(c+1) - \frac{1}{c+1}$$

$$= \ln\frac{c+5}{c+1} + \frac{1}{c+5} - \frac{1}{c+1} = \ln 3 - \frac{1}{3}$$

Comparing: $\frac{c+5}{c+1} = 3 \Rightarrow c+5 = 3c+3 \Rightarrow 2c = 2 \Rightarrow c = 1$

Check: $\frac{1}{1+5} - \frac{1}{1+1} = \frac{1}{6} - \frac{3}{6} = -\frac{2}{6} = -\frac{1}{3}$ ✓ so $c = 1$

6

$$\tan x - \frac{3}{\tan x} = 2 \Rightarrow \tan^2 x - 3 = 2\tan x$$

$$\tan^2 x - 2\tan x - 3 = 0$$

$$(\tan x - 3)(\tan x + 1) = 0 \Rightarrow \tan x = 3 \text{ or } \tan x = -1$$

$\tan x = 3$: $x = 71.565...° = 71.6°$, $x = 180° + 71.6° = 251.6°$

$\tan x = -1$: $x = 135°$, $x = 315°$

$x = 71.6°, 135°, 251.6°, 315°$

7

$$\text{LHS} = \sin 8\theta \tan 4\theta + \cos 8\theta$$

$$= 2\sin 4\theta \cos 4\theta \times \frac{\sin 4\theta}{\cos 4\theta} + \cos 8\theta$$

$$= 2\sin^2 4\theta + \cos 8\theta$$

$$= 2\sin^2 4\theta + 1 - 2\sin^2 4\theta$$

$$= 1 = \text{RHS}$$

8 (a)

$$\cos x + \sqrt{3}\sin x = R\sin(x+\alpha) = R\sin x\cos\alpha + R\cos x\sin\alpha$$

$R\cos\alpha = \sqrt{3}$, $R\sin\alpha = 1$

$R^2 = (\sqrt{3})^2 + 1^2 = 4 \Rightarrow R = 2$

$\tan\alpha = \frac{1}{\sqrt{3}} \Rightarrow \alpha = \frac{\pi}{6}$

$\cos x + \sqrt{3}\sin x = 2\sin\left(x + \frac{\pi}{6}\right)$ $R = 2$, $\alpha = \frac{\pi}{6}$

8 (b)

$$\cos x = \sqrt{3} - \sqrt{3}\sin x \Rightarrow \cos x + \sqrt{3}\sin x = \sqrt{3}$$


$$2\sin\left(x + \frac{\pi}{6}\right) = \sqrt{3} \Rightarrow \sin\left(x + \frac{\pi}{6}\right) = \frac{\sqrt{3}}{2}$$

$$-\pi \leq x \leq \pi \Rightarrow -\frac{5\pi}{6} \leq x + \frac{\pi}{6} \leq \frac{7\pi}{6}$$

Question number	
	$x + \frac{\pi}{6} = \frac{\pi}{3}$ or $\frac{2\pi}{3}$
	$x = \frac{\pi}{6}$ or $x = \frac{\pi}{2}$
9 (a)	$f(1) = 1^4 - 1 - \frac{1}{3-2} = -1 < 0$
	$f(2) = 2^4 - 2 - \frac{1}{6-2} = 13.75 > 0$
	There is a change of sign and f is continuous for $1 \leq x \leq 2$
	(the asymptote is at $x = \frac{2}{3}$), so there is a root between 1 and 2.
9 (b)	$f(1.125) < 0$ and $f(1.25) > 0$, so use the midpoint: $x = 1.1875$
9 (c)	$f(1.15) = -0.0906... < 0$ and $f(1.25) = 0.620 > 0$
	so the root lies between 1.15 and 1.25: root = 1.2 (1 d.p.)
9 (d)(i)	$f(0) = 0.5 > 0$ and $f(1) = -1 < 0$, so there is a change of
	sign between $x = 0$ and $x = 1$.
9 (d)(ii)	$f(x)$ is undefined at $x = \frac{2}{3}$ (vertical asymptote, f is discontinuous).
	The x values are homing in on $\frac{2}{3}$ and the $f(x)$ values are diverging
	(alternating sign, getting larger), so the sign change is caused by the
	asymptote, not a root. Fig. 9.1 shows there is no root between 0 and 1.

Question number

10



$f'(x) = 0$ at $x = 0$ and $x \approx \pm 2.1$ (turning points of f); odd function

11 (a)

$g'(x) = \sin x + x\cos x - \sin x = x\cos x$

Gradient 1: $x\cos x = 1 \Rightarrow \cos x = \frac{1}{x} \Rightarrow \frac{1}{x} - \cos x = 0$

11 (b)

Let $f(x) = \frac{1}{x} - \cos x$, $f'(x) = -\frac{1}{x^2} + \sin x$

Iteration: $x_{n+1} = x_n - \frac{1/x_n - \cos x_n}{-1/x_n^2 + \sin x_n}$

Starting value $x_1 = 5$ (from Fig. 11.2 the smallest positive root is near 5)

$x_2 = 4.916247\ldots$, $x_3 = 4.917185\ldots$, $x_4 = 4.917185\ldots$

$x = 4.9172$ (4 d.p.)

11 (c)

At $x = 3$ the curve $y = f(x)$ is near a turning point, so $f'(3) \approx 0.030$ is close to zero: the tangent is almost horizontal, so the next iteration $x_2 = 3 - \frac{f(3)}{f'(3)} \approx -41.1$ is a long way from the root.

12 (a)

$\frac{dx}{d\theta} = 2\cos 2\theta$, $\frac{dy}{d\theta} = -2\sin\theta - 2\sin 2\theta$

Stationary points: $\frac{dy}{dx} = \frac{dy/d\theta}{dx/d\theta} = 0 \Rightarrow \frac{dy}{d\theta} = 0$

Question number	
	$-2\sin\theta - 2(2\sin\theta\cos\theta) = 0 \Rightarrow -2\sin\theta(1 + 2\cos\theta) = 0$
	$\sin\theta = 0$ or $\cos\theta = -\frac{1}{2}$
	$\theta = 0, \pi$ or $\theta = \frac{2\pi}{3}, \frac{4\pi}{3}$
	$\theta = 0$: $x = 0 + 2 = 2$, $y = 2 + 1 = 3$
	$\theta = \pi$: $x = 0 + 2 = 2$, $y = -2 + 1 = -1$
	$\theta = \frac{2\pi}{3}$: $x = \sin\frac{4\pi}{3} + 2 = 2 - \frac{\sqrt{3}}{2}$
	$y = 2(-\frac{1}{2}) + \cos\frac{4\pi}{3} = -1 - \frac{1}{2} = -\frac{3}{2}$
	$\theta = \frac{4\pi}{3}$: $x = \sin\frac{8\pi}{3} + 2 = 2 + \frac{\sqrt{3}}{2}$
	$y = 2(-\frac{1}{2}) + \cos\frac{8\pi}{3} = -1 - \frac{1}{2} = -\frac{3}{2}$
	$\frac{dx}{d\theta} = 2, 2, -1, -1$ at these values ($\neq 0$), so all are stationary points.
	Stationary points: $(2, 3)$, $(2, -1)$, $\left(2 - \frac{\sqrt{3}}{2}, -\frac{3}{2}\right)$, $\left(2 + \frac{\sqrt{3}}{2}, -\frac{3}{2}\right)$
12 (b)	$x = 2$
13	A $(-3, 9)$: $t_1 = -3$; B $(1, 1)$: $t_2 = 1$
	$t_1t_2 = -3 \times 1 = -3$, which is the y-coordinate of the intersection $(-1, -3)$ ✓
14	$t_1 = -3$, $t_2 = 1$:
	$t_1^2 + t_2^2 + t_1t_2 + \frac{1}{2} = (-3)^2 + 1^2 + (-3)(1) + \frac{1}{2} = 9 + 1 - 3 + 0.5 = 7.5$
	which is the y-coordinate of the intersection $(-12, 7.5)$ ✓
15 (a)	$\frac{dy}{dx} = 2ax + b$, so at $x = t$ the gradient is $2at + b$
	and the point is $(t, at^2 + bt + c)$
	$y - (at^2 + bt + c) = (2at + b)(x - t)$
	$y = (2at + b)x - 2at^2 - bt + at^2 + bt + c$
	$y = (2at + b)x - at^2 + c$ as required
15 (b)	Tangents at P and Q: $y = (2ax_P + b)x - ax_P^2 + c$, $y = (2ax_Q + b)x - ax_Q^2 + c$
	They cross where $(2ax_P + b)x - ax_P^2 + c = (2ax_Q + b)x - ax_Q^2 + c$

Question number

$(2ax_P - 2ax_Q)x = ax_P^2 - ax_Q^2$

$x = \frac{a(x_P^2 - x_Q^2)}{2a(x_P - x_Q)} = \frac{(x_P + x_Q)(x_P - x_Q)}{2(x_P - x_Q)} = \frac{x_P + x_Q}{2} \quad (x_P \neq x_Q)$

which is the mean of the x-coordinates of P and Q.

16 $a\left(\frac{x_P + x_Q}{2}\right)^2 + b\left(\frac{x_P + x_Q}{2}\right) + c - a\left(\frac{x_P - x_Q}{2}\right)^2$

$= a\left(\frac{x_P^2 + 2x_Px_Q + x_Q^2}{4} - \frac{x_P^2 - 2x_Px_Q + x_Q^2}{4}\right) + b\left(\frac{x_P + x_Q}{2}\right) + c$

$= a\left(\frac{4x_Px_Q}{4}\right) + b\left(\frac{x_P + x_Q}{2}\right) + c$

$= ax_Px_Q + b\left(\frac{x_P + x_Q}{2}\right) + c$ as required

17 $y = x^2 \Rightarrow \frac{dy}{dx} = 2x$, so the gradient of the tangent at (t, t^2) is $2t$

Gradient of normal $= -\frac{1}{2t}$

$y - t^2 = -\frac{1}{2t}(x - t) \Rightarrow y = -\frac{x}{2t} + \frac{t}{2t} + t^2$

$y = -\frac{x}{2t} + t^2 + \frac{1}{2}$ as required

18 From line 19: $y = ax^2 + bx + c - \frac{ah^2}{4}$ with $a = 6, b = -7, c = 1, h = 5$

$y = 6x^2 - 7x + 1 - \frac{6 \times 5^2}{4} = 6x^2 - 7x + 1 - 37.5$

$y = 6x^2 - 7x - 36.5$