OCR
Oxford Cambridge and RSA

Tuesday 14 June 2022 – Afternoon

A Level Mathematics A

H240/02 Pure Mathematics and Statistics

Time allowed: 2 hours

6 0 8 6 2 0 0 6 2 8

You must have:
- the Printed Answer Booklet
- a scientific or graphical calculator

INSTRUCTIONS
- Use black ink. You can use an HB pencil, but only for graphs and diagrams.
- Write your answer to each question in the space provided in the **Printed Answer Booklet**. If you need extra space use the lined pages at the end of the Printed Answer Booklet. The question numbers must be clearly shown.
- Fill in the boxes on the front of the Printed Answer Booklet.
- Answer **all** the questions.
- Where appropriate, your answer should be supported with working. Marks might be given for using a correct method, even if your answer is wrong.
- Give non-exact numerical answers correct to **3** significant figures unless a different degree of accuracy is specified in the question.
- The acceleration due to gravity is denoted by $g\,\text{m}\,\text{s}^{-2}$. When a numerical value is needed use $g = 9.8$ unless a different value is specified in the question.
- Do **not** send this Question Paper for marking. Keep it in the centre or recycle it.

INFORMATION
- The total mark for this paper is **100**.
- The marks for each question are shown in brackets **[ ]**.
- This document has **12** pages.

ADVICE
- Read each question carefully before you start your answer.

© OCR 2022 [603/1038/8]
DC (LK/FC) 306682/4

OCR is an exempt Charity

stemup.co.uk

Formulae
A Level Mathematics A (H240)

Arithmetic series

$S_n = \frac{1}{2}n(a+l) = \frac{1}{2}n\{2a+(n-1)d\}$

Geometric series

$S_n = \dfrac{a(1-r^n)}{1-r}$

$S_\infty = \dfrac{a}{1-r}$ for $|r| < 1$

Binomial series

$(a+b)^n = a^n + {}^n\mathrm{C}_1 a^{n-1}b + {}^n\mathrm{C}_2 a^{n-2}b^2 + \ldots + {}^n\mathrm{C}_r a^{n-r}b^r + \ldots + b^n \qquad (n \in \mathbb{N}),$

where ${}^n\mathrm{C}_r = {}_n\mathrm{C}_r = \binom{n}{r} = \dfrac{n!}{r!(n-r)!}$

$(1+x)^n = 1 + nx + \dfrac{n(n-1)}{2!}x^2 + \ldots + \dfrac{n(n-1)\ldots(n-r+1)}{r!}x^r + \ldots \qquad (|x| < 1, n \in \mathbb{R})$

Differentiation

$\mathrm{f}(x)$	$\mathrm{f}'(x)$
$\tan kx$	$k\sec^2 kx$
$\sec x$	$\sec x \tan x$
$\cot x$	$-\mathrm{cosec}^2 x$
$\mathrm{cosec}\, x$	$-\mathrm{cosec}\, x \cot x$

Quotient rule $y = \dfrac{u}{v}, \dfrac{\mathrm{d}y}{\mathrm{d}x} = \dfrac{v\frac{\mathrm{d}u}{\mathrm{d}x} - u\frac{\mathrm{d}v}{\mathrm{d}x}}{v^2}$

Differentiation from first principles

$\mathrm{f}'(x) = \lim_{h \to 0} \dfrac{\mathrm{f}(x+h) - \mathrm{f}(x)}{h}$

Integration

$\displaystyle\int \frac{\mathrm{f}'(x)}{\mathrm{f}(x)}\,\mathrm{d}x = \ln|\mathrm{f}(x)| + c$

$\displaystyle\int \mathrm{f}'(x)\big(\mathrm{f}(x)\big)^n\,\mathrm{d}x = \frac{1}{n+1}\big(\mathrm{f}(x)\big)^{n+1} + c$

Integration by parts $\displaystyle\int u\frac{\mathrm{d}v}{\mathrm{d}x}\,\mathrm{d}x = uv - \int v\frac{\mathrm{d}u}{\mathrm{d}x}\,\mathrm{d}x$

Small angle approximations

$\sin\theta \approx \theta$, $\cos\theta \approx 1 - \frac{1}{2}\theta^2$, $\tan\theta \approx \theta$ where θ is measured in radians

© OCR 2022

stemup.co.uk

Trigonometric identities

$\sin(A \pm B) = \sin A \cos B \pm \cos A \sin B$

$\cos(A \pm B) = \cos A \cos B \mp \sin A \sin B$

$\tan(A \pm B) = \dfrac{\tan A \pm \tan B}{1 \mp \tan A \tan B} \qquad \left(A \pm B \neq (k + \frac{1}{2})\pi\right)$

Numerical methods

Trapezium rule: $\displaystyle\int_a^b y\,\mathrm{d}x \approx \tfrac{1}{2}h\{(y_0 + y_n) + 2(y_1 + y_2 + \ldots + y_{n-1})\}$, where $h = \dfrac{b-a}{n}$

The Newton-Raphson iteration for solving $\mathrm{f}(x) = 0$: $x_{n+1} = x_n - \dfrac{\mathrm{f}(x_n)}{\mathrm{f}'(x_n)}$

Probability

$\mathrm{P}(A \cup B) = \mathrm{P}(A) + \mathrm{P}(B) - \mathrm{P}(A \cap B)$

$\mathrm{P}(A \cap B) = \mathrm{P}(A)\mathrm{P}(B \mid A) = \mathrm{P}(B)\mathrm{P}(A \mid B)$ **or** $\mathrm{P}(A \mid B) = \dfrac{\mathrm{P}(A \cap B)}{\mathrm{P}(B)}$

Standard deviation

$\sqrt{\dfrac{\Sigma(x-\overline{x})^2}{n}} = \sqrt{\dfrac{\Sigma x^2}{n} - \overline{x}^2}$ **or** $\sqrt{\dfrac{\Sigma f(x-\overline{x})^2}{\Sigma f}} = \sqrt{\dfrac{\Sigma f x^2}{\Sigma f} - \overline{x}^2}$

The binomial distribution

If $X \sim B(n, p)$ then $P(X = x) = \dbinom{n}{x} p^x (1-p)^{n-x}$, mean of X is np, variance of X is $np(1-p)$

Hypothesis test for the mean of a normal distribution

If $X \sim \mathrm{N}(\mu, \sigma^2)$ then $\overline{X} \sim \mathrm{N}\left(\mu, \dfrac{\sigma^2}{n}\right)$ and $\dfrac{\overline{X} - \mu}{\sigma/\sqrt{n}} \sim \mathrm{N}(0,1)$

Percentage points of the normal distribution

If Z has a normal distribution with mean 0 and variance 1 then, for each value of p, the table gives the value of z such that $P(Z \leqslant z) = p$.

p	0.75	0.90	0.95	0.975	0.99	0.995	0.9975	0.999	0.9995
z	0.674	1.282	1.645	1.960	2.326	2.576	2.807	3.090	3.291

Kinematics

Motion in a straight line	Motion in two dimensions
$v = u + at$	$\mathbf{v} = \mathbf{u} + \mathbf{a}t$
$s = ut + \frac{1}{2}at^2$	$\mathbf{s} = \mathbf{u}t + \frac{1}{2}\mathbf{a}t^2$
$s = \frac{1}{2}(u+v)t$	$\mathbf{s} = \frac{1}{2}(\mathbf{u}+\mathbf{v})t$
$v^2 = u^2 + 2as$	
$s = vt - \frac{1}{2}at^2$	$\mathbf{s} = \mathbf{v}t - \frac{1}{2}\mathbf{a}t^2$

stemup.co.uk

Section A: Pure Mathematics

Answer **all** the questions.

1 **In this question you must show detailed reasoning.**

Solve the following equations.

(a) $\frac{x}{x+1} - \frac{x-1}{x+2} = 0$ **[3]**

(b) $\frac{8}{x^6} - \frac{7}{x^3} - 1 = 0$ **[3]**

(c) $3^{x^2-7} = \frac{1}{243}$ **[2]**

2 The points A and B have position vectors $3\mathbf{i} + 2\mathbf{j}$ and $4\mathbf{i} + 2\mathbf{j} - 5\mathbf{k}$ respectively.

(a) Find the length of AB. **[2]**

Point P has position vector $p\mathbf{i} - 3\mathbf{k}$, where p is a constant. P lies on the circumference of a circle of which AB is a diameter.

(b) Find the two possible values of p. **[3]**

© OCR 2022

stemup.co.uk

3 **(a)** Amaya and Ben integrated $(1+x)^2$, with respect to x, using different methods, as follows.

Amaya: $\int (1+x)^2 \,\mathrm{d}x = \dfrac{(1+x)^3}{3} + c \qquad = \frac{1}{3} + x + x^2 + \frac{1}{3}x^3 + c$

Ben: $\int (1+x)^2 \,\mathrm{d}x = \int (1+2x+x^2) \,\mathrm{d}x \qquad = x + x^2 + \frac{1}{3}x^3 + c$

Charlie said that, because these answers are different, at least one of them must be wrong.

Explain whether you agree with Charlie's statement. **[1]**

(b) You are given that a is a constant greater than 1.

(i) Find $\int_1^a \dfrac{1}{(1+x)^2} \,\mathrm{d}x$, giving your answer as a single fraction in terms of the constant a. **[3]**

(ii) You are given that the area enclosed by the curve $y = \dfrac{1}{(1+x)^2}$, the x-axis and the lines $x = 1$ and $x = a$ is equal to $\frac{1}{3}$.

Determine the value of a. **[2]**

(c) **In this question you must show detailed reasoning.**

Find the exact value of $\int_0^{\frac{1}{12}\pi} \dfrac{\cos 2x}{\sin 2x + 2} \,\mathrm{d}x$, giving your answer in its simplest form. **[4]**

4 An artist is creating a design for a large painting. The design includes a set of steps of varying heights. In the painting the lowest step has height 20 cm and the height of each other step is 5% less than the height of the step immediately below it.

In the painting the total height of the steps is 205 cm, correct to the nearest centimetre.

Determine the number of steps in the design. **[5]**

5 **In this question you must show detailed reasoning.**

A curve has equation $y = x^3 - 3x^2 + 4x$.

(a) Show that the curve has no stationary points. **[2]**

(b) Show that the curve has exactly one point of inflection. **[2]**

stemup.co.uk

6 **(a)** The diagrams show five different graphs. In each case the whole of the graph is shown.




Fig. 1.1




Fig. 1.2




Fig. 1.3




Fig. 1.4




Fig. 1.5

Place ticks in the boxes in the table in the Printed Answer Booklet to indicate, for each graph, whether it represents a one-one function, a many-one function, a function that is its own inverse or it does not represent a function. There may be more than one tick in some rows or columns of the table. **[4]**

(b) A function f is defined by $\mathrm{f}(x) = \frac{1}{x}$ for the domain $\{x : 0 < x \leqslant 2\}$.

State the range of f, giving your answer in set notation. **[2]**

© OCR 2022

stemup.co.uk

7 It is given that any integer can be expressed in the form $3m+r$, where m is an integer and r is 0, 1 or 2.


Use this fact to answer the following.

(a) By considering the different values of r, prove that the square of any integer **cannot** be expressed in the form $3n+2$, where n is an integer. **[4]**

(b) Three integers are chosen at random from the integers 1 to 99 inclusive. The three integers are not necessarily different.

By considering the different values of r, determine the probability that the sum of these three integers is divisible by 3. **[4]**

8



The diagram shows a water tank which is shaped as an inverted cone with semi-vertical angle 30° and height 50 cm. Initially the tank is full, and the depth of the water is 50 cm.

Water flows out of a small hole at the bottom of the tank. The rate at which the water flows out is modelled by $\frac{dV}{dt} = -2h$, where $V\,\text{cm}^3$ is the volume of water remaining and h cm is the depth of water in the tank t seconds after the water begins to flow out.

Determine the time taken for the tank to become empty.


[For a cone with base radius r and height h the volume V is given by $\frac{1}{3}\pi r^2 h$.] **[7]**

stemup.co.uk

Section B: Statistics

Answer **all** the questions.

9 The heights, in centimetres, of a random sample of 150 plants of a certain variety were measured. The results are summarised in the histogram.



One of the 150 plants is chosen at random, and its height, X cm, is noted.

(a) Show that $P(20 < X < 30) = 0.147$, correct to **3** significant figures. **[2]**

Sam suggests that the distribution of X can be well modelled by the distribution N(40, 100).

(b) (i) Give a brief justification for the use of the normal distribution in this context. **[1]**

(ii) Give a brief justification for the choice of the parameter values 40 and 100. **[2]**

(c) Use Sam's model to find $P(20 < X < 30)$. **[1]**

Nina suggests a different model. She uses the midpoints of the classes to calculate estimates, m and s, for the mean and standard deviation respectively, in centimetres, of the 150 heights. She then uses the distribution $N(m, s^2)$ as her model.

(d) Use Nina's model to find $P(20 < X < 30)$. **[4]**

(e) (i) Complete the table in the Printed Answer Booklet to show the probabilities obtained from Sam's model and Nina's model. **[2]**

(ii) By considering the different ranges of values of X given in the table, discuss how well the two models fit the original distribution. **[2]**

© OCR 2022

stemup.co.uk

BLANK PAGE

Turn over for question 10

stemup.co.uk

10 The table shows the age structure of usual residents of 18 Local Authorities (LAs) in the North West region of the UK in 2011.

Local Authority	Age 0 to 17	Age 18 to 24	Age 25 to 64	Age 65 and over
A	26.20%	9.06%	51.81%	12.92%
B	23.32%	8.99%	52.32%	15.37%
C	22.24%	8.96%	52.56%	16.23%
D	22.67%	8.10%	53.27%	15.96%
E	20.70%	7.77%	54.77%	16.76%
F	18.14%	6.51%	51.13%	24.21%
G	18.96%	14.20%	48.51%	18.33%
H	19.06%	14.79%	52.12%	14.04%
I	25.15%	9.04%	51.16%	14.65%
J	22.93%	8.81%	52.22%	16.04%
K	21.48%	13.98%	50.82%	13.73%
L	23.98%	9.20%	52.26%	14.56%
M	21.67%	11.19%	52.94%	14.19%
N	17.82%	6.01%	51.93%	24.23%
O	22.83%	7.30%	53.86%	16.01%
P	21.76%	8.28%	54.03%	15.93%
Q	21.42%	8.43%	53.90%	16.25%
R	18.61%	7.33%	49.35%	24.71%

Percentage of residents

(a) Without reference to any other columns, explain how you would use **only** the columns for the age ranges 0 to 17 and 18 to 24 to decide whether an LA might be one of the following.

(i) An LA that includes a university **[1]**

(ii) An LA that attracts young couples to live **[1]**

(iii) An LA that attracts retired people to live **[1]**

(b) Using your answers to part **(a)**, identify the following.

(i) Four LAs that might include a university **[1]**

(ii) Three LAs that might be attractive to retired people **[1]**

(c) Explain why your answer to part **(b)(ii)**, based only on the columns for the age ranges 0 to 17 and 18 to 24, may not be reliable. **[1]**

© OCR 2022

stemup.co.uk

(d) The lower quartile, median and upper quartile of the percentages in the column "Age 65 and over" are 14.56%, 15.99% and 16.76% respectively.

Use this information to comment on your answers to part **(b)(ii)** and part **(c)**. **[2]**

In a magazine article, a councillor plans to describe a typical LA in the North West region. He wants to quote the average percentage of residents aged 65 or over.

(e) The mean of the percentages in the column "Age 65 and over" is 16.90%.

Use this information, and the information given in part **(d)**, to explain whether the median or the mean better represents the data in the column "Age 65 and over". **[2]**

11 In the past the masses of new-born babies in a certain country were normally distributed with mean 3300 g. Last year a publicity campaign was held to encourage pregnant women to improve their diet.

Following this campaign, it is required to test whether the mean mass of new-born babies has increased. A random sample of 200 new-born babies is chosen, and it is found that their mean mass is 3360 g. It is given that the standard deviation of the masses of new-born babies is 450 g.

Carry out the test at the 2.5% significance level. **[7]**

12 A firm claims that no more than 2% of their packets of sugar are underweight. A market researcher believes that the actual proportion is greater than 2%. In order to test the firm's claim, the researcher weighs a random sample of 600 packets and carries out a hypothesis test, at the 5% significance level, using the null hypothesis $p = 0.02$.

(a) Given that the researcher's null hypothesis is correct, determine the probability that the researcher will conclude that the firm's claim is incorrect. **[5]**

(b) The researcher finds that 18 out of the 600 packets are underweight. A colleague says

"18 out of 600 is 3%, so there is evidence that the actual proportion of underweight bags is greater than 2%."

Criticise this statement. **[1]**

Turn over for question 13

stemup.co.uk

13 There are 25 students in a class.

- The number of students who study both History and English is 3.
- The number of students who study neither History nor English is 14.
- The number of students who study History but not English is three times the number who study English but not History.

(a)
- Show this information on a Venn diagram.
- Determine the probability that a student selected at random studies English. **[4]**

Two different students from the class are chosen at random.

(b) Given that exactly one of the two students studies English, determine the probability that exactly one of the two students studies History. **[6]**

END OF QUESTION PAPER

OCR
Oxford Cambridge and RSA

Copyright Information

OCR is committed to seeking permission to reproduce all third-party content that it uses in its assessment materials. OCR has attempted to identify and contact all copyright holders whose work is used in this paper. To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced in the OCR Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download from our public website (www.ocr.org.uk) after the live examination series.

If OCR has unwittingly failed to correctly acknowledge or clear any third-party content in this assessment material, OCR will be happy to correct its mistake at the earliest possible opportunity.

For queries or further information please contact The OCR Copyright Team, The Triangle Building, Shaftesbury Road, Cambridge CB2 8EA.

OCR is part of Cambridge University Press & Assessment, which is itself a department of the University of Cambridge.

© OCR 2022

stemup.co.uk

Surname	StemUp Model Solution
Answer booklet	Write the question number in the left-hand margin at the start of each answer.

Question number

1 (a)

$x(x + 2) - (x - 1)(x + 1) = 0$ (multiplying by $(x + 1)(x + 2)$)

$x^2 + 2x - (x^2 - 1) = 0$

$2x + 1 = 0$

$x = -\frac{1}{2}$

1 (b)

Multiply by x^6: $8 - 7x^3 - x^6 = 0 \Rightarrow x^6 + 7x^3 - 8 = 0$

Let $u = x^3$: $u^2 + 7u - 8 = 0$

$(u + 8)(u - 1) = 0 \Rightarrow u = -8$ or $u = 1$

$x^3 = -8 \Rightarrow x = -2$; $x^3 = 1 \Rightarrow x = 1$

$x = 1$ or $x = -2$

1 (c)

$3^{x^2 - 7} = 1/243 = 3^{-5}$ (since $243 = 3^5$)

$x^2 - 7 = -5 \Rightarrow x^2 = 2$

$x = \pm\sqrt{2}$

2 (a)

$AB = (4i + 2j - 5k) - (3i + 2j) = i - 5k$

$|AB| = \sqrt{(1^2 + 0^2 + (-5)^2)} = \sqrt{26}$ (= 5.10 to 3 s.f.)

2 (b)

P lies on the circle with diameter AB, so angle APB = 90° $\Rightarrow$ PA · PB = 0

$PA = (3i + 2j) - (pi - 3k) = (3 - p)i + 2j + 3k$

$PB = (4i + 2j - 5k) - (pi - 3k) = (4 - p)i + 2j - 2k$

$PA \cdot PB = (3 - p)(4 - p) + 4 - 6 = 0$

$p^2 - 7p + 12 - 2 = 0 \Rightarrow p^2 - 7p + 10 = 0$

$(p - 2)(p - 5) = 0 \Rightarrow p = 2$ or $p = 5$

Question number	
3 (a)	No, I disagree with Charlie. The two answers differ only by the constant ⅓, and ⅓ can be absorbed into the arbitrary constant of integration (Ben's c = Amaya's c + ⅓), so both answers are correct.
3 (b)(i)	$\int_1^a (1+x)^{-2}\,dx = [-(1+x)^{-1}]_1^a = -\frac{1}{1+a} + \frac{1}{2}$ $= \frac{(1+a)-2}{2(1+a)} = \frac{a-1}{2(a+1)}$
3 (b)(ii)	$\frac{a-1}{2(a+1)} = \frac{1}{3} \Rightarrow 3(a-1) = 2(a+1)$ $3a - 3 = 2a + 2 \Rightarrow a = 5$
3 (c)	$\int_0^{\pi/12} \frac{\cos 2x}{\sin 2x + 2}\,dx = [\tfrac{1}{2}\ln\lvert \sin 2x + 2 \rvert]_0^{\pi/12}$ (since d/dx (sin 2x + 2) = 2 cos 2x) $= \tfrac{1}{2}(\ln(\sin(\pi/6) + 2) - \ln(\sin 0 + 2))$ $= \tfrac{1}{2}(\ln\frac{5}{2} - \ln 2) = \tfrac{1}{2}\ln\frac{5}{4}$
4	Heights form a GP with a = 20, r = 0.95 $S_n = \frac{20(1-0.95^n)}{1-0.95} = 400(1-0.95^n) = 205$ $1 - 0.95^n = \frac{205}{400} \Rightarrow 0.95^n = \frac{195}{400} = 0.4875$ $n = \frac{\ln 0.4875}{\ln 0.95} = 14.006\ldots \approx 14$ Check: $S_{14} = 400(1-0.95^{14}) = 204.93\ldots = 205$ cm (nearest cm) Number of steps = 14

Question number

5 (a) $y = x^3 - 3x^2 + 4x \Rightarrow dy/dx = 3x^2 - 6x + 4$

Stationary points where $3x^2 - 6x + 4 = 0$

Discriminant $b^2 - 4ac = (-6)^2 - 4 \times 3 \times 4 = 36 - 48 = -12 < 0$

so $dy/dx = 0$ has no real roots, hence the curve has no stationary points.

5 (b) $d^2y/dx^2 = 6x - 6 = 0 \Rightarrow x = 1$ (the only root)

For $x < 1$, $d^2y/dx^2 < 0$ and for $x > 1$, $d^2y/dx^2 > 0$, so d^2y/dx^2 changes sign at $x = 1$: $x = 1$ gives a point of inflection, and since $6x - 6 = 0$ has only one root, the curve has exactly one point of inflection.

6 (a)

	One-one	Many-one	Own inverse	Not a function
Fig. 1.1		✓		
Fig. 1.2				✓
Fig. 1.3	✓		✓	
Fig. 1.4	✓			
Fig. 1.5		✓		

6 (b) f is decreasing on $0 < x \leq 2$, $f(2) = \frac{1}{2}$ and $f(x) \to \infty$ as $x \to 0$

Range: $\{ y : y \geq \frac{1}{2} \}$

7 (a) Any integer is $3m$, $3m + 1$ or $3m + 2$.

$r = 0$: $(3m)^2 = 9m^2 = 3(3m^2)$, of the form $3n$

$r = 1$: $(3m + 1)^2 = 9m^2 + 6m + 1 = 3(3m^2 + 2m) + 1$, of the form $3n + 1$

$r = 2$: $(3m + 2)^2 = 9m^2 + 12m + 4 = 3(3m^2 + 4m + 1) + 1$, form $3n + 1$

None of these is of the form $3n + 2$, so the square of any integer cannot be expressed in the form $3n + 2$.

Question number	
7 (b)	1 to 99 contains 33 integers with each of r = 0, 1, 2,
	so P(r = 0) = P(r = 1) = P(r = 2) = ⅓
	The sum is divisible by 3 if the three remainders are all the same, or all different.
	All the same: $P(0,0,0) + P(1,1,1) + P(2,2,2) = 3 \times (\frac{1}{3})^3 = 3/27$
	All different (0, 1, 2 in any order): $3! \times (\frac{1}{3})^3 = 6/27$
	P(sum divisible by 3) = 3/27 + 6/27 = 9/27 = ⅓
8	Radius of water surface $r = h \tan 30° = h/\sqrt{3}$
	$V = \frac{1}{3}\pi\left(\frac{h}{\sqrt{3}}\right)^2 h = \frac{\pi}{9}h^3 \Rightarrow \frac{dV}{dh} = \frac{\pi}{3}h^2$
	$\frac{dV}{dt} = \frac{dV}{dh} \times \frac{dh}{dt} = \frac{\pi}{3}h^2\frac{dh}{dt} = -2h$
	$\Rightarrow \pi h \frac{dh}{dt} = -6 \Rightarrow \int_{50}^{0} \pi h\, dh = \int_0^T -6\, dt$
	$\left[\frac{\pi h^2}{2}\right]_{50}^{0} = \left[-6t\right]_0^T \Rightarrow 0 - \frac{\pi \times 50^2}{2} = -6T$
	$T = \frac{2500\pi}{12} = \frac{625\pi}{3} = 654.49...$
	Time taken for the tank to empty = 625π/3 s ≈ 654 seconds (3 s.f.)
9 (a)	Using small squares: area of 20 to 30 block = 110 small squares,
	total area of histogram = 750 small squares
	P(20 < X < 30) = 110/750 = 0.14666... = 0.147 (3 s.f.)
9 (b)(i)	The histogram is roughly bell-shaped: roughly symmetrical with a single peak in
	the middle, tailing off at each end.
9 (b)(ii)	The distribution is roughly symmetrical about 40 (the peak is at about 40), so μ
	≈ 40. Almost all the data lie between 10 and 70, i.e. within about 30 of 40, and

Question number

$70 - 40 \approx 3\sigma$, so $\sigma \approx 10$, giving $\sigma^2 = 100$.

9 (c)

$X \sim N(40, 100)$: $P(20 < X < 30) = 0.13590... = 0.136$ (3 s.f.)

9 (d)

Midpoints 15, 25, 32.5, 37.5, 42.5, 47.5, 55, 65

Frequencies 4, 22, 23, 28, 29, 22, 20, 2 ($\Sigma f = 150$)

$\Sigma fx = 5915$, $m = 5915/150 = 39.433... = 39.4$

$\Sigma fx^2 = 249287.5$, $s^2 = 249287.5/150 - 39.433...^2 = 106.93...$

$s = 10.340... = 10.3$

$X \sim N(39.433, 10.341^2)$: $P(20 < X < 30) = 0.15071... = 0.151$ (3 s.f.)

9 (e)(i)

x	< 20	20 to 30	30 to 35	35 to 40	40 to 45	45 to 50	50 to 60	> 60
Histogram	0.027	0.147	0.153	0.187	0.193	0.147	0.133	0.013
N(40, 100)	0.023	0.136	0.150	0.191	0.191	0.150	0.136	0.023
$N(m, s^2)$	0.030	0.151	0.153	0.189	0.183	0.142	0.130	0.023

9 (e)(ii)

Both models are fairly close to the histogram. For the lower values of X (below 40) Nina's model is the better fit, e.g. for 20 to 30 Nina gives 0.151 against 0.147 while Sam gives 0.136. For the higher values of X (above 40) Sam's model is the better fit, e.g. for 40 to 45 Sam gives 0.191 against 0.193 while Nina gives 0.183.

10 (a)(i)

An LA with a university would have a high percentage of residents aged 18 to 24.

10 (a)(ii)

An LA attracting young couples would have a high percentage aged 0 to 17 (children) and a high percentage aged 18 to 24.

Question number	
10 (a)(iii)	An LA attracting retired people would have a low percentage aged 0 to 17 and a low percentage aged 18 to 24.
10 (b)(i)	G, H, K, M
10 (b)(ii)	F, N, R
10 (c)	A low percentage of residents aged 0 to 17 and 18 to 24 does not necessarily mean a high percentage aged 65 and over: the LA might instead have a large percentage aged 25 to 64, so the other age ranges need to be considered.
10 (d)	IQR = 16.76 − 14.56 = 2.20 UQ + 1.5 × IQR = 16.76 + 1.5 × 2.20 = 20.06 F (24.21%), N (24.23%) and R (24.71%) are all greater than 20.06, so all three are outliers with an unusually high percentage aged 65 and over. This confirms that F, N and R are likely to be attractive to retired people, despite the concern in part (c).
10 (e)	The mean (16.90%) is greater than the upper quartile (16.76%), because it is pulled up by the three high outliers F, N and R. So the median (15.99%) better represents the data in the column "Age 65 and over".
11	Let μ be the population mean mass of new-born babies (in grams). H_0: $\mu = 3300$ H_1: $\mu > 3300$ Under H_0, $\bar{X} \sim N(3300, \frac{450^2}{200}) = N(3300, 1012.5)$ Sample mean $\bar{x} = 3360$ $P(\bar{X} \geq 3360) = 0.029673... = 0.0297$ (3 s.f.) $0.0297 > 0.025$, so the result is not significant: do not reject H_0.

Question number

There is insufficient evidence at the 2.5% level to suggest that the mean mass of new-born babies has increased.

12 (a) Let X = number of underweight packets in the sample: $X \sim B(600, 0.02)$

H_0: $p = 0.02$, H_1: $p > 0.02$, one-tailed test at 5%

$P(X \geqslant 18) = 1 - P(X \leqslant 17) = 0.0610$ (3 s.f.) > 0.05

$P(X \geqslant 19) = 1 - P(X \leqslant 18) = 0.0359$ (3 s.f.) < 0.05

So the critical region is $X \geqslant 19$.


P(researcher concludes the claim is incorrect | H_0 true) $= P(X \geqslant 19)$

$= 0.03588... = 0.0359$ (3 s.f.)

12 (b) The statement is wrong: the critical region is $X \geqslant 19$, so 18 is in the acceptance region ($P(X \geqslant 18) = 0.0610 > 0.05$). At the 5% level there is insufficient evidence that the proportion is greater than 2%; a sample proportion above 2% is not enough on its own, a proper hypothesis test is needed.

13 (a) Let x = number studying English but not History, so History only = $3x$.

$3x + 3 + x + 14 = 25 \Rightarrow 4x = 8 \Rightarrow x = 2$



H = History
E = English

Number studying English = $2 + 3 = 5$

P(English) = 5/25 = 1/5

Question number

13 (b)

Not English: 6 + 14 = 20 students. (Choosing 2 without replacement)

$P(\text{exactly one English}) = \frac{5}{25} \times \frac{20}{24} \times 2 = \frac{200}{600} = \frac{1}{3}$

Exactly one English and exactly one History:

either (English only and History only) or (both and neither)

$P = \left(\frac{2}{25} \times \frac{6}{24} + \frac{3}{25} \times \frac{14}{24}\right) \times 2 = \frac{(12 + 42) \times 2}{600} = \frac{108}{600} = \frac{9}{50}$

$P(\text{exactly one History} \mid \text{exactly one English}) = \frac{9/50}{1/3} = \frac{27}{50} \quad (= 0.54)$