OCR
Oxford Cambridge and RSA

Wednesday 07 October 2020 – Afternoon

A Level Mathematics A

H240/01 Pure Mathematics

Time allowed: 2 hours

8232526068

You must have:
- the Printed Answer Booklet
- a scientific or graphical calculator

INSTRUCTIONS
- Use black ink. You can use an HB pencil, but only for graphs and diagrams.
- Write your answer to each question in the space provided in the **Printed Answer Booklet**. If you need extra space use the lined pages at the end of the Printed Answer Booklet. The question numbers must be clearly shown.
- Fill in the boxes on the front of the Printed Answer Booklet.
- Answer **all** the questions.
- Where appropriate, your answer should be supported with working. Marks might be given for using a correct method, even if your answer is wrong.
- Give non-exact numerical answers correct to 3 significant figures unless a different degree of accuracy is specified in the question.
- The acceleration due to gravity is denoted by $g\,\mathrm{m\,s^{-2}}$. When a numerical value is needed use $g = 9.8$ unless a different value is specified in the question.
- Do **not** send this Question Paper for marking. Keep it in the centre or recycle it.

INFORMATION
- The total mark for this paper is **100**.
- The marks for each question are shown in brackets **[ ]**.
- This document has **8** pages.

ADVICE
- Read each question carefully before you start your answer.

© OCR 2020 [603/1038/8]
DC (ST/CT) 193126/7

OCR is an exempt Charity

stemup.co.uk

Formulae
A Level Mathematics A (H240)

Arithmetic series

$S_n = \frac{1}{2}n(a+l) = \frac{1}{2}n\{2a+(n-1)d\}$

Geometric series

$$S_n = \frac{a(1-r^n)}{1-r}$$

$$S_\infty = \frac{a}{1-r} \quad \text{for } |r| < 1$$

Binomial series

$$(a+b)^n = a^n + {}^n\mathrm{C}_1 a^{n-1}b + {}^n\mathrm{C}_2 a^{n-2}b^2 + \ldots + {}^n\mathrm{C}_r a^{n-r}b^r + \ldots + b^n \qquad (n \in \mathbb{N})$$

where ${}^n\mathrm{C}_r = {}_n\mathrm{C}_r = \binom{n}{r} = \frac{n!}{r!(n-r)!}$

$$(1+x)^n = 1 + nx + \frac{n(n-1)}{2!}x^2 + \ldots + \frac{n(n-1)\ldots(n-r+1)}{r!}x^r + \ldots \qquad (|x| < 1, n \in \mathbb{R})$$

Differentiation

$\mathrm{f}(x)$	$\mathrm{f}'(x)$
$\tan kx$	$k\sec^2 kx$
$\sec x$	$\sec x \tan x$
$\cot x$	$-\mathrm{cosec}^2 x$
$\mathrm{cosec}\, x$	$-\mathrm{cosec}\, x \cot x$

Quotient rule $y = \frac{u}{v}, \frac{\mathrm{d}y}{\mathrm{d}x} = \frac{v\frac{\mathrm{d}u}{\mathrm{d}x} - u\frac{\mathrm{d}v}{\mathrm{d}x}}{v^2}$

Differentiation from first principles

$$\mathrm{f}'(x) = \lim_{h \to 0} \frac{\mathrm{f}(x+h) - \mathrm{f}(x)}{h}$$

Integration

$$\int \frac{\mathrm{f}'(x)}{\mathrm{f}(x)}\,\mathrm{d}x = \ln|\mathrm{f}(x)| + c$$

$$\int \mathrm{f}'(x)\big(\mathrm{f}(x)\big)^n\,\mathrm{d}x = \frac{1}{n+1}\big(\mathrm{f}(x)\big)^{n+1} + c$$

Integration by parts $\int u\frac{\mathrm{d}v}{\mathrm{d}x}\,\mathrm{d}x = uv - \int v\frac{\mathrm{d}u}{\mathrm{d}x}\,\mathrm{d}x$

Small angle approximations

$\sin\theta \approx \theta$, $\cos\theta \approx 1 - \frac{1}{2}\theta^2$, $\tan\theta \approx \theta$ where θ is measured in radians

© OCR 2020

stemup.co.uk

Trigonometric identities

$\sin(A \pm B) = \sin A \cos B \pm \cos A \sin B$

$\cos(A \pm B) = \cos A \cos B \mp \sin A \sin B$

$\tan(A \pm B) = \dfrac{\tan A \pm \tan B}{1 \mp \tan A \tan B} \qquad \left(A \pm B \neq (k + \frac{1}{2})\pi\right)$

Numerical methods

Trapezium rule: $\displaystyle\int_a^b y\,\mathrm{d}x \approx \tfrac{1}{2}h\{(y_0 + y_n) + 2(y_1 + y_2 + \ldots + y_{n-1})\}$, where $h = \dfrac{b-a}{n}$

The Newton-Raphson iteration for solving $\mathrm{f}(x) = 0$: $x_{n+1} = x_n - \dfrac{\mathrm{f}(x_n)}{\mathrm{f}'(x_n)}$

Probability

$\mathrm{P}(A \cup B) = \mathrm{P}(A) + \mathrm{P}(B) - \mathrm{P}(A \cap B)$

$\mathrm{P}(A \cap B) = \mathrm{P}(A)\mathrm{P}(B \,|\, A) = \mathrm{P}(B)\mathrm{P}(A \,|\, B)$ **or** $\mathrm{P}(A \,|\, B) = \dfrac{\mathrm{P}(A \cap B)}{\mathrm{P}(B)}$

Standard deviation

$\sqrt{\dfrac{\Sigma(x-\bar{x})^2}{n}} = \sqrt{\dfrac{\Sigma x^2}{n} - \bar{x}^2}$ **or** $\sqrt{\dfrac{\Sigma f(x-\bar{x})^2}{\Sigma f}} = \sqrt{\dfrac{\Sigma f x^2}{\Sigma f} - \bar{x}^2}$

The binomial distribution

If $X \sim \mathrm{B}(n, p)$ then $\mathrm{P}(X = x) = \dbinom{n}{x} p^x (1-p)^{n-x}$, mean of X is np, variance of X is $np(1-p)$

Hypothesis test for the mean of a normal distribution

If $X \sim \mathrm{N}(\mu, \sigma^2)$ then $\overline{X} \sim \mathrm{N}\left(\mu, \dfrac{\sigma^2}{n}\right)$ and $\dfrac{\overline{X} - \mu}{\sigma/\sqrt{n}} \sim \mathrm{N}(0, 1)$

Percentage points of the normal distribution

If Z has a normal distribution with mean 0 and variance 1 then, for each value of p, the table gives the value of z such that $\mathrm{P}(Z \leqslant z) = p$.

p	0.75	0.90	0.95	0.975	0.99	0.995	0.9975	0.999	0.9995
z	0.674	1.282	1.645	1.960	2.326	2.576	2.807	3.090	3.291

Kinematics

Motion in a straight line	Motion in two dimensions
$v = u + at$	$\mathbf{v} = \mathbf{u} + \mathbf{a}t$
$s = ut + \frac{1}{2}at^2$	$\mathbf{s} = \mathbf{u}t + \frac{1}{2}\mathbf{a}t^2$
$s = \frac{1}{2}(u + v)t$	$\mathbf{s} = \frac{1}{2}(\mathbf{u} + \mathbf{v})t$
$v^2 = u^2 + 2as$	
$s = vt - \frac{1}{2}at^2$	$\mathbf{s} = \mathbf{v}t - \frac{1}{2}\mathbf{a}t^2$

© OCR 2020

stemup.co.uk

Answer **all** the questions.

1 **(a)** For a small angle θ, where θ is in radians, show that $2\cos\theta + (1-\tan\theta)^2 \approx 3 - 2\theta$. **[3]**

(b) Hence determine an approximate solution to $2\cos\theta + (1-\tan\theta)^2 = 28\sin\theta$. **[2]**

2 Simplify fully.

(a) $\sqrt{12a} \times \sqrt{3a^5}$ **[2]**

(b) $(64b^3)^{\frac{1}{3}} \times (4b^4)^{-\frac{1}{2}}$ **[2]**

(c) $7 \times 9^{3c} - 4 \times 27^{2c}$ **[4]**

3 A cylindrical metal tin of radius r cm is closed at both ends. It has a volume of $16000\pi\,\text{cm}^3$.

(a) Show that its total surface area, $A\,\text{cm}^2$, is given by $A = 2\pi r^2 + 32000\pi r^{-1}$. **[4]**


(b) Use calculus to determine the minimum total surface area of the tin. You should justify that it is a minimum. **[6]**

4 Prove by contradiction that there is no greatest multiple of 5. **[3]**

© OCR 2020

stemup.co.uk

5



The diagram shows points A and B, which have position vectors **a** and **b** with respect to an origin O. P is the point on OB such that $OP : PB = 3:1$ and Q is the midpoint of AB.

(a) Find $\overrightarrow{PQ}$ in terms of **a** and **b**. **[2]**

The line OA is extended to a point R, so that PQR is a straight line.

(b) Explain why $\overrightarrow{PR} = k(2\mathbf{a} - \mathbf{b})$, where k is a constant. **[2]**

(c) Hence determine the ratio $OA : AR$. **[4]**

6 A mobile phone company records their annual sales on 31st December every year.

Paul thinks that the annual sales, S million, can be modelled by the equation $S = ab^t$, where a and b are both positive constants and t is the number of years since 31st December 2015.

Paul tests his theory by using the annual sales figures from 31st December 2015 to 31st December 2019. He plots these results on a graph, with t on the horizontal axis and $\log_{10} S$ on the vertical axis.

(a) Explain why, if Paul's model is correct, the results should lie on a straight line of best fit on his graph. **[3]**

The results lie on a straight line of best fit which has a gradient of 0.146 and an intercept on the vertical axis of 0.583.

(b) Use these values to obtain estimates for a and b, correct to 2 significant figures. **[2]**

(c) Use this model to predict the year in which, on the 31st December, the annual sales would first be recorded as greater than 200 million. **[3]**

(d) Give a reason why this prediction may not be reliable. **[1]**

stemup.co.uk

7 Two students, Anna and Ben, are starting a revision programme. They will both revise for 30 minutes on Day 1. Anna will increase her revision time by 15 minutes for every subsequent day. Ben will increase his revision time by 10% for every subsequent day.

(a) Verify that on Day 10 Anna does 94 minutes more revision than Ben, correct to the nearest minute. **[3]**

Let Day X be the first day on which Ben does more revision than Anna.

(b) Show that X satisfies the inequality $X > \log_{1.1}(0.5X + 0.5) + 1$. **[3]**

(c) Use the iterative formula $x_{n+1} = \log_{1.1}(0.5x_n + 0.5) + 1$ with $x_1 = 10$ to find the value of X.

You should show the result of each iteration. **[3]**


(d) (i) Give a reason why Anna's revision programme may not be realistic. **[1]**

(ii) Give a **different** reason why Ben's revision programme may not be realistic. **[1]**

8 **(a)** Differentiate $(2 + 3x^2)e^{2x}$ with respect to x. **[3]**

(b) Hence show that $(2 + 3x^2)e^{2x}$ is increasing for all values of x. **[4]**

9



The diagram shows the graph of $y = |2x - 3|$.

(a) State the coordinates of the points of intersection with the axes. **[2]**

(b) Given that the graphs of $y = |2x - 3|$ and $y = ax + 2$ have two distinct points of intersection, determine


(i) the set of possible values of a, **[4]**

(ii) the x-coordinates of the points of intersection of these graphs, giving your answers in terms of a. **[3]**

© OCR 2020

stemup.co.uk

10



The diagram shows the curve $y = \sin\left(\frac{1}{2}\sqrt{x-1}\right)$, for $1 \leqslant x \leqslant 2$.

(a) Use rectangles of width 0.25 to find upper and lower bounds for $\int_1^2 \sin\left(\frac{1}{2}\sqrt{x-1}\right)\mathrm{d}x$. Give your answers correct to 3 significant figures. **[4]**


(b) (i) Use the substitution $t = \sqrt{x-1}$ to show that $\int \sin\left(\frac{1}{2}\sqrt{x-1}\right)\mathrm{d}x = \int 2t\sin\left(\frac{1}{2}t\right)\mathrm{d}t$. **[3]**

(ii) Hence show that $\int_1^2 \sin\left(\frac{1}{2}\sqrt{x-1}\right)\mathrm{d}x = 8\sin\frac{1}{2} - 4\cos\frac{1}{2}$. **[4]**

Turn over for questions 11 and 12

stemup.co.uk

11 In this question you must show detailed reasoning.



The diagram shows a circle with equation $x^2 + y^2 - 10x - 14y + 64 = 0$. A tangent is drawn from the point $P(0, 2)$ to meet the circle at the point A. The equation of this tangent is of the form $y = mx + 2$, where m is a constant **greater than** 1.

(a) (i) Show that the x-coordinate of A satisfies the equation $(m^2 + 1)x^2 - 10(m + 1)x + 40 = 0$. **[2]**

(ii) Hence determine the equation of the tangent to the circle at A which passes through P. **[4]**

A second tangent is drawn from P to meet the circle at a second point B. The equation of this tangent is of the form $y = nx + 2$, where n is a constant **less than** 1.

(b) Determine the exact value of $\tan APB$. **[4]**

12 Find the general solution of the differential equation

$$(2x^3 - 3x^2 - 11x + 6)\frac{dy}{dx} = y(20x - 35).$$

Give your answer in the form $y = f(x)$. **[9]**

END OF QUESTION PAPER

OCR
Oxford Cambridge and RSA

Copyright Information

OCR is committed to seeking permission to reproduce all third-party content that it uses in its assessment materials. OCR has attempted to identify and contact all copyright holders whose work is used in this paper. To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced in the OCR Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download from our public website (www.ocr.org.uk) after the live examination series.

If OCR has unwittingly failed to correctly acknowledge or clear any third-party content in this assessment material, OCR will be happy to correct its mistake at the earliest possible opportunity.

For queries or further information please contact The OCR Copyright Team, The Triangle Building, Shaftesbury Road, Cambridge CB2 8EA.

OCR is part of the Cambridge Assessment Group; Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.

© OCR 2020

stemup.co.uk

Surname	StemUp Model Solution
Answer booklet	Write the question number in the left-hand margin at the start of each answer.

Question number

1 (a)

For small θ: $\cos\theta \approx 1 - \frac{1}{2}\theta^2$, $\tan\theta \approx \theta$

$2\cos\theta + (1 - \tan\theta)^2 \approx 2(1 - \frac{1}{2}\theta^2) + (1 - \theta)^2$

$= 2 - \theta^2 + 1 - 2\theta + \theta^2$

$= 3 - 2\theta$ as required

1 (b)

$\sin\theta \approx \theta$, so $3 - 2\theta = 28\theta$

$30\theta = 3$

$\theta = 0.1$

2 (a)

$\sqrt{(12a)} \times \sqrt{(3a^5)} = \sqrt{(36a^6)}$

$= 6a^3$

2 (b)

$(64b^3)^{1/3} = 4b$, $(4b^4)^{-1/2} = \frac{1}{\sqrt{(4b^8)}} = \frac{1}{2b^2}$

$4b \times \frac{1}{2b^2} = \frac{2}{b}$

2 (c)

$9^{3c} = (3^2)^{3c} = 3^{6c}$

$27^{2c} = (3^3)^{2c} = 3^{6c}$

$7 \times 9^{3c} - 4 \times 27^{2c} = 7 \times 3^{6c} - 4 \times 3^{6c}$

$= 3 \times 3^{6c}$

$= 3^{6c+1}$

Question number

3 (a)

Volume: $\pi r^2 h = 16000\pi$

$h = \frac{16000}{r^2} = 16000r^{-2}$

Surface area = two circular ends + curved surface

$A = 2\pi r^2 + 2\pi rh$

$= 2\pi r^2 + 2\pi r \times 16000r^{-2}$

$= 2\pi r^2 + 32000\pi r^{-1}$ as required

3 (b)

$\frac{dA}{dr} = 4\pi r - 32000\pi r^{-2}$

At a stationary point $\frac{dA}{dr} = 0$: $4\pi r - 32000\pi r^{-2} = 0$

$4\pi r^3 = 32000\pi \Rightarrow r^3 = 8000 \Rightarrow r = 20$

$A = 2\pi(20)^2 + 32000\pi(20)^{-1} = 800\pi + 1600\pi = 2400\pi$

$\frac{d^2A}{dr^2} = 4\pi + 64000\pi r^{-3}$

When $r = 20$: $\frac{d^2A}{dr^2} = 4\pi + 64000\pi \div 8000 = 12\pi \ (= 37.7) > 0$

so this is a minimum.

Minimum surface area = 2400π cm^2 $(= 7540$ cm^2 to 3 s.f.$)$

4

Assume there is a greatest multiple of 5, N, so $N = 5k$ for some integer k.

Consider $N + 5 = 5k + 5 = 5(k + 1)$.

$k + 1$ is an integer, so $N + 5$ is a multiple of 5, and $N + 5 > N$.

This contradicts the assumption that N is the greatest multiple of 5.

Hence there is no greatest multiple of 5.

Question number

5 (a) $\overrightarrow{OP} = \frac{3}{4}\underline{b}$, $\overrightarrow{PB} = \frac{1}{4}\underline{b}$, $\overrightarrow{BQ} = \frac{1}{2}\overrightarrow{BA} = \frac{1}{2}(\underline{a} - \underline{b})$

$\overrightarrow{PQ} = \overrightarrow{PB} + \overrightarrow{BQ} = \frac{1}{4}\underline{b} + \frac{1}{2}(\underline{a} - \underline{b}) = \frac{1}{2}\underline{a} - \frac{1}{4}\underline{b}$

5 (b) P, Q and R lie on a straight line, so $\overrightarrow{PR}$ is in the same direction as $\overrightarrow{PQ}$ so $\overrightarrow{PR}$ is a multiple of $\overrightarrow{PQ}$:

$\overrightarrow{PR} = \lambda(\frac{1}{2}\underline{a} - \frac{1}{4}\underline{b}) = \frac{\lambda}{4}(2\underline{a} - \underline{b})$

so $\overrightarrow{PR} = k(2\underline{a} - \underline{b})$ where $k = \frac{\lambda}{4}$ is a constant

5 (c) $\overrightarrow{AR} = \overrightarrow{AO} + \overrightarrow{OP} + \overrightarrow{PR} = -\underline{a} + \frac{3}{4}\underline{b} + k(2\underline{a} - \underline{b})$

$= (2k - 1)\underline{a} + (\frac{3}{4} - k)\underline{b}$

R lies on OA extended, so $\overrightarrow{AR}$ is a multiple of $\underline{a}$ only:

$\frac{3}{4} - k = 0 \Rightarrow k = \frac{3}{4}$

$\overrightarrow{AR} = (\frac{3}{2} - 1)\underline{a} = \frac{1}{2}\underline{a}$, $\overrightarrow{OA} = \underline{a}$

$OA : AR = 1 : \frac{1}{2} = 2 : 1$

Question number

6 (a)

$S = ab^t$

$\log_{10} S = \log_{10}(ab^t) = \log_{10} a + \log_{10} b^t$

$\log_{10} S = (\log_{10} b)\, t + \log_{10} a$

This is of the form $Y = mX + c$ with $Y = \log_{10} S$ and $X = t$,

gradient $m = \log_{10} b$ and intercept $c = \log_{10} a$ (both constants),

so the graph of $\log_{10} S$ against t is a straight line.

6 (b)

$\log_{10} a = 0.583 \Rightarrow a = 10^{0.583} = 3.828\ldots = 3.8$ (2 s.f.)

$\log_{10} b = 0.146 \Rightarrow b = 10^{0.146} = 1.399\ldots = 1.4$ (2 s.f.)

6 (c)

$3.8 \times 1.4^t > 200$

$1.4^t > \dfrac{200}{3.8} = 52.63\ldots$

$t > \dfrac{\log 52.63\ldots}{\log 1.4} = 11.779\ldots$

so $t = 12$ (first whole number of years after 31st December 2015)

2015 + 12 = 2027: the year is 2027

6 (d)

It is extrapolating far beyond the data (2015 to 2019); sales are unlikely

to keep growing at the same rate as the market is limited by the number of customers.

7 (a)

Anna (AP, $a = 30$, $d = 15$): $u_{10} = 30 + 9 \times 15 = 165$ minutes

Ben (GP, $a = 30$, $r = 1.1$): $u_{10} = 30 \times 1.1^9 = 70.738\ldots$ minutes

$165 - 70.738\ldots = 94.26\ldots$ = 94 minutes (nearest minute), as required

Question number	
7 (b)	Anna: $u_X = 30 + 15(X - 1)$, Ben: $u_X = 30 \times 1.1^{X-1}$
	Ben does more than Anna: $30 \times 1.1^{X-1} > 30 + 15(X - 1)$
	$30 \times 1.1^{X-1} > 15X + 15$
	$1.1^{X-1} > 0.5X + 0.5$
	Taking $\log_{1.1}$ (an increasing function): $X - 1 > \log_{1.1}(0.5X + 0.5)$
	$X > \log_{1.1}(0.5X + 0.5) + 1$ as required
7 (c)	$x_1 = 10$
	$x_2 = \log_{1.1}(0.5 \times 10 + 0.5) + 1 = 18.886...$
	$x_3 = 25.099...$, $x_4 = 27.951...$, $x_5 = 29.039...$
	$x_6 = 29.426...$, $x_7 = 29.561...$, $x_8 = 29.607...$
	$x_9 = 29.623...$, $x_{10} = 29.628...$ → converging to 29.63...
	X is the first whole number greater than 29.63..., so $X = 30$
7 (d)(i)	Her time keeps increasing: by Day 96 she would need more than 24 hours
	of revision in a day, so there will not be enough hours in the day.
7 (d)(ii)	Increasing by 10% gives times with decimal parts of a minute (e.g. 33 min,
	36.3 min, 39.93 min), which cannot realistically be timed that accurately.
8 (a)	Let $u = 2 + 3x^2$, $v = e^{2x}$: $\frac{du}{dx} = 6x$, $\frac{dv}{dx} = 2e^{2x}$
	$\frac{d}{dx}[(2 + 3x^2)e^{2x}] = 6xe^{2x} + (2 + 3x^2)(2e^{2x})$
	$= e^{2x}(6x^2 + 6x + 4)$

Question number	
8 (b)	$e^{2x} > 0$ for all x
	$6x^2 + 6x + 4 = 6(x^2 + x) + 4 = 6(x + \frac{1}{2})^2 - \frac{3}{2} + 4 = 6(x + \frac{1}{2})^2 + \frac{5}{2}$
	$6(x + \frac{1}{2})^2 \geq 0$, so the minimum value is $\frac{5}{2} > 0$, so $6x^2 + 6x + 4 > 0$ for all x
	Hence the gradient $e^{2x}(6x^2 + 6x + 4) > 0$ for all x,
	so $(2 + 3x^2)e^{2x}$ is increasing for all values of x.
9 (a)	$y = 0$: $2x - 3 = 0 \Rightarrow x = 1.5$ $\quad$ $(1.5, 0)$
	$x = 0$: $y = \lvert -3 \rvert = 3$ $\quad$ $(0, 3)$
9 (b)(i)	$y = ax + 2$ passes through $(0, 2)$, which lies below the graph ($\lvert 0 - 3 \rvert = 3 > 2$).
	For two intersections the line must meet both arms of the graph.
	It must meet the right arm (gradient 2), so $a < 2$.
	Boundary: the line passes through the vertex $(1.5, 0)$:
	$0 = 1.5a + 2 \Rightarrow a = -\frac{4}{3}$
	For $a \leq -\frac{4}{3}$ the line meets the graph at most once, so need $a > -\frac{4}{3}$
	$-\frac{4}{3} < a < 2$
9 (b)(ii)	Right arm: $2x - 3 = ax + 2 \Rightarrow (2 - a)x = 5 \Rightarrow x = \frac{5}{2 - a}$
	Left arm: $3 - 2x = ax + 2 \Rightarrow (2 + a)x = 1 \Rightarrow x = \frac{1}{2 + a}$

Question number

10 (a)

x:	1	1.25	1.5	1.75	2
y:	0	0.24740	0.34623	0.41961	0.47943

The curve is increasing, so left-hand heights give the lower bound:

Lower $= 0.25\{\sin 0 + \sin(\tfrac{1}{2}\sqrt{0.25}) + \sin(\tfrac{1}{2}\sqrt{0.5}) + \sin(\tfrac{1}{2}\sqrt{0.75})\}$

$= 0.25 \times 1.01325\ldots = 0.25331\ldots = 0.253$

Upper $= 0.25\{\sin(\tfrac{1}{2}\sqrt{0.25}) + \sin(\tfrac{1}{2}\sqrt{0.5}) + \sin(\tfrac{1}{2}\sqrt{0.75}) + \sin(\tfrac{1}{2}\sqrt{1})\}$

$= 0.25 \times 1.49267\ldots = 0.37317\ldots = 0.373$

$0.253 <$ integral < 0.373

10 (b)(i)

$t = \sqrt{(x-1)} \Rightarrow t^2 = x - 1$

$2t\,\dfrac{dt}{dx} = 1 \Rightarrow dx = 2t\,dt$

$\int \sin(\tfrac{1}{2}\sqrt{(x-1)})\,dx = \int \sin(\tfrac{1}{2}t) \times 2t\,dt = \int 2t\sin(\tfrac{1}{2}t)\,dt$ as required

10 (b)(ii)

Limits: $x = 1 \Rightarrow t = 0$, $x = 2 \Rightarrow t = 1$

By parts: $u = 2t$, $\dfrac{dv}{dt} = \sin(\tfrac{1}{2}t) \Rightarrow \dfrac{du}{dt} = 2$, $v = -2\cos(\tfrac{1}{2}t)$

$\int 2t\sin(\tfrac{1}{2}t)\,dt = -4t\cos(\tfrac{1}{2}t) + \int 4\cos(\tfrac{1}{2}t)\,dt$

$= -4t\cos(\tfrac{1}{2}t) + 8\sin(\tfrac{1}{2}t)$

$\displaystyle\int_0^1 2t\sin(\tfrac{1}{2}t)\,dt = [-4t\cos(\tfrac{1}{2}t) + 8\sin(\tfrac{1}{2}t)]$ from $t = 0$ to $t = 1$

$= (-4\cos\tfrac{1}{2} + 8\sin\tfrac{1}{2}) - (-0 + 8\sin 0)$

$= (-4\cos\tfrac{1}{2} + 8\sin\tfrac{1}{2}) - 0$

So $\displaystyle\int_1^2 \sin(\tfrac{1}{2}\sqrt{(x-1)})\,dx = 8\sin\frac{1}{2} - 4\cos\frac{1}{2}$ as required

Question number

11 (a)(i) Substitute $y = mx + 2$ into $x^2 + y^2 - 10x - 14y + 64 = 0$:

$x^2 + (mx + 2)^2 - 10x - 14(mx + 2) + 64 = 0$

$x^2 + m^2x^2 + 4mx + 4 - 10x - 14mx - 28 + 64 = 0$

$(m^2 + 1)x^2 - 10(m + 1)x + 40 = 0$ as required

11 (a)(ii) Tangent, so the quadratic has one repeated root: $b^2 - 4ac = 0$

$100(m + 1)^2 - 4 \times 40(m^2 + 1) = 0$

$100m^2 + 200m + 100 - 160m^2 - 160 = 0$

$-60m^2 + 200m - 60 = 0 \quad \Rightarrow \quad 3m^2 - 10m + 3 = 0$

$(3m - 1)(m - 3) = 0$

$m = 3$ or $m = \frac{1}{3}$; $m > 1$ so $m = 3$

Tangent: $y = 3x + 2$

11 (b) $(x - 5)^2 + (y - 7)^2 = 25 + 49 - 64 = 10$: centre $C(5, 7)$, radius $\sqrt{10}$

$PC^2 = (5 - 0)^2 + (7 - 2)^2 = 50$, so $PC = 5\sqrt{2}$

Radius $CA \perp$ tangent PA: $PA^2 = PC^2 - CA^2 = 50 - 10 = 40$, $PA = 2\sqrt{10}$

In right-angled triangle PAC: $\tan APC = \frac{CA}{PA} = \frac{\sqrt{10}}{2\sqrt{10}} = \frac{1}{2}$

By symmetry about PC, angle $APB = 2 \times$ angle APC

$\tan APB = \frac{2\tan APC}{1 - \tan^2 APC} = \frac{2 \times \frac{1}{2}}{1 - \frac{1}{4}} = \frac{1}{\frac{3}{4}} = \frac{4}{3}$

Question number

12

$$\int \frac{1}{y}\,dy = \int \frac{20x - 35}{2x^3 - 3x^2 - 11x + 6}\,dx$$

Let $f(x) = 2x^3 - 3x^2 - 11x + 6$: $f(3) = 54 - 27 - 33 + 6 = 0$

so $(x - 3)$ is a factor of $f(x)$

$2x^3 - 3x^2 - 11x + 6 = (x - 3)(2x^2 + 3x - 2) = (x - 3)(x + 2)(2x - 1)$

$$\frac{20x - 35}{(x + 2)(x - 3)(2x - 1)} \equiv \frac{A}{x + 2} + \frac{B}{x - 3} + \frac{C}{2x - 1}$$

$20x - 35 \equiv A(x - 3)(2x - 1) + B(x + 2)(2x - 1) + C(x + 2)(x - 3)$

$x = -2$: $-75 = 25A \Rightarrow A = -3$ $\quad$ $x = 3$: $25 = 25B \Rightarrow B = 1$

$x = \frac{1}{2}$: $-25 = C(\frac{5}{2})(-\frac{5}{2}) \Rightarrow C = 4$

$$\int \frac{1}{y}\,dy = \int \left(-\frac{3}{x + 2} + \frac{1}{x - 3} + \frac{4}{2x - 1}\right) dx$$

$\ln|y| = -3\ln|x + 2| + \ln|x - 3| + 2\ln|2x - 1| + \ln A$

$$\ln|y| = \ln \frac{A(x - 3)(2x - 1)^2}{(x + 2)^3}$$

$$y = \frac{A(x - 3)(2x - 1)^2}{(x + 2)^3}$$ where A is a constant