OCR
Oxford Cambridge and RSA

Wednesday 6 October 2021 – Afternoon

A Level Mathematics A

H240/01 Pure Mathematics

Time allowed: 2 hours

8947468997

You must have:
- the Printed Answer Booklet
- a scientific or graphical calculator

INSTRUCTIONS
- Use black ink. You can use an HB pencil, but only for graphs and diagrams.
- Write your answer to each question in the space provided in the **Printed Answer Booklet**. If you need extra space use the lined pages at the end of the Printed Answer Booklet. The question numbers must be clearly shown.
- Fill in the boxes on the front of the Printed Answer Booklet.
- Answer **all** the questions.
- Where appropriate, your answer should be supported with working. Marks might be given for using a correct method, even if your answer is wrong.
- Give non-exact numerical answers correct to **3** significant figures unless a different degree of accuracy is specified in the question.
- The acceleration due to gravity is denoted by $g\,\text{m}\,\text{s}^{-2}$. When a numerical value is needed use $g = 9.8$ unless a different value is specified in the question.
- Do **not** send this Question Paper for marking. Keep it in the centre or recycle it.

INFORMATION
- The total mark for this paper is **100**.
- The marks for each question are shown in brackets **[ ]**.
- This document has **8** pages.

ADVICE
- Read each question carefully before you start your answer.

© OCR 2021 [603/1038/8]
DC (KN/FC) 300260/3
OCR is an exempt Charity

stemup.co.uk

Formulae
A Level Mathematics A (H240)

Arithmetic series

$S_n = \frac{1}{2}n(a+l) = \frac{1}{2}n\{2a+(n-1)d\}$

Geometric series

$S_n = \dfrac{a(1-r^n)}{1-r}$

$S_\infty = \dfrac{a}{1-r}$ for $|r| < 1$

Binomial series

$(a+b)^n = a^n + {}^n\mathrm{C}_1 a^{n-1}b + {}^n\mathrm{C}_2 a^{n-2}b^2 + \ldots + {}^n\mathrm{C}_r a^{n-r}b^r + \ldots + b^n \qquad (n \in \mathbb{N})$

where ${}^n\mathrm{C}_r = {}_n\mathrm{C}_r = \binom{n}{r} = \dfrac{n!}{r!(n-r)!}$

$(1+x)^n = 1 + nx + \dfrac{n(n-1)}{2!}x^2 + \ldots + \dfrac{n(n-1)\ldots(n-r+1)}{r!}x^r + \ldots \qquad (|x| < 1, n \in \mathbb{R})$

Differentiation

$\mathrm{f}(x)$	$\mathrm{f}'(x)$
$\tan kx$	$k\sec^2 kx$
$\sec x$	$\sec x \tan x$
$\cot x$	$-\mathrm{cosec}^2 x$
$\mathrm{cosec}\, x$	$-\mathrm{cosec}\, x \cot x$

Quotient rule $y = \dfrac{u}{v}, \dfrac{\mathrm{d}y}{\mathrm{d}x} = \dfrac{v\frac{\mathrm{d}u}{\mathrm{d}x} - u\frac{\mathrm{d}v}{\mathrm{d}x}}{v^2}$

Differentiation from first principles

$\mathrm{f}'(x) = \lim_{h \to 0} \dfrac{\mathrm{f}(x+h) - \mathrm{f}(x)}{h}$

Integration

$\displaystyle\int \frac{\mathrm{f}'(x)}{\mathrm{f}(x)}\,\mathrm{d}x = \ln|\mathrm{f}(x)| + c$

$\displaystyle\int \mathrm{f}'(x)\big(\mathrm{f}(x)\big)^n\,\mathrm{d}x = \frac{1}{n+1}\big(\mathrm{f}(x)\big)^{n+1} + c$

Integration by parts $\displaystyle\int u\frac{\mathrm{d}v}{\mathrm{d}x}\,\mathrm{d}x = uv - \int v\frac{\mathrm{d}u}{\mathrm{d}x}\,\mathrm{d}x$

Small angle approximations

$\sin\theta \approx \theta$, $\cos\theta \approx 1 - \frac{1}{2}\theta^2$, $\tan\theta \approx \theta$ where θ is measured in radians

© OCR 2021

stemup.co.uk

Trigonometric identities

$\sin(A \pm B) = \sin A \cos B \pm \cos A \sin B$

$\cos(A \pm B) = \cos A \cos B \mp \sin A \sin B$

$\tan(A \pm B) = \dfrac{\tan A \pm \tan B}{1 \mp \tan A \tan B} \qquad \left(A \pm B \neq (k + \frac{1}{2})\pi\right)$

Numerical methods

Trapezium rule: $\displaystyle\int_a^b y\,\mathrm{d}x \approx \tfrac{1}{2}h\{(y_0 + y_n) + 2(y_1 + y_2 + \ldots + y_{n-1})\}$, where $h = \dfrac{b-a}{n}$

The Newton-Raphson iteration for solving $\mathrm{f}(x) = 0$: $x_{n+1} = x_n - \dfrac{\mathrm{f}(x_n)}{\mathrm{f}'(x_n)}$

Probability

$\mathrm{P}(A \cup B) = \mathrm{P}(A) + \mathrm{P}(B) - \mathrm{P}(A \cap B)$

$\mathrm{P}(A \cap B) = \mathrm{P}(A)\mathrm{P}(B \mid A) = \mathrm{P}(B)\mathrm{P}(A \mid B)$ **or** $\mathrm{P}(A \mid B) = \dfrac{\mathrm{P}(A \cap B)}{\mathrm{P}(B)}$

Standard deviation

$\sqrt{\dfrac{\Sigma(x-\bar{x})^2}{n}} = \sqrt{\dfrac{\Sigma x^2}{n} - \bar{x}^2}$ **or** $\sqrt{\dfrac{\Sigma f(x-\bar{x})^2}{\Sigma f}} = \sqrt{\dfrac{\Sigma f x^2}{\Sigma f} - \bar{x}^2}$

The binomial distribution

If $X \sim \mathrm{B}(n, p)$ then $\mathrm{P}(X = x) = \dbinom{n}{x} p^x (1-p)^{n-x}$, mean of X is np, variance of X is $np(1-p)$

Hypothesis test for the mean of a normal distribution

If $X \sim \mathrm{N}(\mu, \sigma^2)$ then $\overline{X} \sim \mathrm{N}\left(\mu, \dfrac{\sigma^2}{n}\right)$ and $\dfrac{\overline{X} - \mu}{\sigma/\sqrt{n}} \sim \mathrm{N}(0, 1)$

Percentage points of the normal distribution

If Z has a normal distribution with mean 0 and variance 1 then, for each value of p, the table gives the value of z such that $\mathrm{P}(Z \leqslant z) = p$.

p	0.75	0.90	0.95	0.975	0.99	0.995	0.9975	0.999	0.9995
z	0.674	1.282	1.645	1.960	2.326	2.576	2.807	3.090	3.291

Kinematics

Motion in a straight line	Motion in two dimensions
$v = u + at$	$\mathbf{v} = \mathbf{u} + \mathbf{a}t$
$s = ut + \frac{1}{2}at^2$	$\mathbf{s} = \mathbf{u}t + \frac{1}{2}\mathbf{a}t^2$
$s = \frac{1}{2}(u + v)t$	$\mathbf{s} = \frac{1}{2}(\mathbf{u} + \mathbf{v})t$
$v^2 = u^2 + 2as$	
$s = vt - \frac{1}{2}at^2$	$\mathbf{s} = \mathbf{v}t - \frac{1}{2}\mathbf{a}t^2$

© OCR 2021

stemup.co.uk

Answer **all** the questions.

1 Determine the set of values of k such that the equation $x^2 + 4x + (k+3) = 0$ has **two** distinct real roots. **[4]**

2 Alex is comparing the cost of mobile phone contracts. Contract ***A*** has a set-up cost of £40 and then costs 4p per minute. Contract ***B*** has no set-up cost, does not charge for the first 100 minutes and then costs 6p per minute.

(a) Find an expression for the cost of each of the contracts in terms of m, where m is the number of minutes for which the phone is used and $m > 100$. **[2]**

(b) Hence find the value of m for which both contracts would cost the same. **[2]**

3 It is given that x is proportional to the product of the square of y and the positive square root of z. When $y = 2$ and $z = 9$, $x = 30$.

(a) Write an equation for x in terms of y and z. **[2]**

(b) Find the value of x when $y = 3$ and $z = 25$. **[2]**

4 In this question you must show detailed reasoning.

The cubic polynomial $\mathrm{f}(x)$ is defined by $\mathrm{f}(x) = 2x^3 - 3x^2 - 11x + 6$.

(a) Use the factor theorem to show that $(2x-1)$ is a factor of $\mathrm{f}(x)$. **[1]**

(b) Express $\mathrm{f}(x)$ in fully factorised form. **[3]**

(c) Hence solve the equation $2 \times 8^y - 3 \times 4^y - 11 \times 2^y + 6 = 0$. **[3]**

© OCR 2021

stemup.co.uk


5 **(a)** The graph of the function $y = \mathrm{f}(x)$ passes through the point P with coordinates (2, 6), and is a one-one function. State the coordinates of the point corresponding to P on each of the following curves.

(i) $y = \mathrm{f}(x) + 3$ **[1]**

(ii) $y = 2\mathrm{f}(3x - 1)$ **[2]**

(iii) $y = \mathrm{f}^{-1}(x)$ **[1]**

(b)



The diagram shows part of the graph of $y = \mathrm{g}'(x)$. This is the graph of the gradient function of $y = \mathrm{g}(x)$. The graph intersects the x-axis at $x = -2$ and $x = 4$.

(i) State the x-coordinate of any stationary points on the graph of $y = \mathrm{g}(x)$. **[1]**

(ii) State the set of values of x for which $y = \mathrm{g}(x)$ is a decreasing function. **[1]**

(iii) State the x-coordinate of any points of inflection on the graph of $y = \mathrm{g}(x)$. **[1]**

6 **(a)** Find the first three terms in the expansion of $(8 - 3x)^{\frac{1}{3}}$ in ascending powers of x. **[4]**

(b) State the range of values of x for which the expansion in part **(a)** is valid. **[1]**

(c) Find the coefficient of x^2 in the expansion of $\dfrac{(8 - 3x)^{\frac{1}{3}}}{(1 + 2x)^2}$. **[4]**

stemup.co.uk

7 The curve $y = (x^2 - 2)\ln x$ has one stationary point which is close to $x = 1$.

(a) Show that the x-coordinate of this stationary point satisfies the equation $2x^2 \ln x + x^2 - 2 = 0$. **[2]**

(b) Show that the Newton-Raphson iterative formula for finding the root of the equation in part **(a)** can be written in the form $x_{n+1} = \dfrac{2x_n^2 \ln x_n + 3x_n^2 + 2}{4x_n(\ln x_n + 1)}$. **[4]**

(c) Apply the Newton-Raphson formula with initial value $x_1 = 1$ to find x_2 and x_3. **[1]**

(d) Find the coordinates of this stationary point, giving each coordinate correct to **3** decimal places. **[2]**

8 Functions f and g are defined for $0 \leqslant x \leqslant 2\pi$ by $\mathrm{f}(x) = 2\tan x$ and $\mathrm{g}(x) = \sec x$.

(a) (i) State the range of f. **[1]**

(ii) State the range of g. **[1]**

(b) (i) Show that $\mathrm{fg}(0.6) = 5.33$, correct to **3** significant figures. **[2]**

(ii) Explain why $\mathrm{f}^{-1}\mathrm{g}(0.6)$ is not defined. **[1]**

(c) In this question you must show detailed reasoning.

Solve the equation $(\mathrm{f}(x))^2 + 6\,\mathrm{g}(x) = 0$. **[5]**

9 A particle moves in the x-y plane so that at time t seconds, where $t \geqslant 0$, its coordinates are given by $x = \mathrm{e}^{2t} - 4\mathrm{e}^{t} + 3$, $y = 2\mathrm{e}^{-3t}$.

(a) Explain why the path of the particle never crosses the x-axis. **[1]**

(b) Determine the exact values of t when the path of the particle intersects the y-axis. **[2]**


(c) Show that $\dfrac{\mathrm{d}y}{\mathrm{d}x} = \dfrac{3}{2\mathrm{e}^{4t} - \mathrm{e}^{5t}}$. **[4]**

(d) Hence find the coordinates of the particle when its path is parallel to the y-axis. **[3]**

© OCR 2021

stemup.co.uk

10 (a)



The diagram shows triangle ABC. The perpendicular from C to AB meets AB at D.

Angle $ACD = x$, angle $DCB = y$, length $BC = a$ and length $AC = b$.

(i) Explain why the length of CD can be written as $a\cos y$. **[1]**


(ii) Show that the area of the triangle ADC is given by $\frac{1}{2}ab\sin x\cos y$. **[1]**

(iii) Hence, or otherwise, show that $\sin(x+y) = \sin x\cos y + \cos x\sin y$. **[4]**

(b) Given that $\sin(30° + \alpha) = \cos(45° - \alpha)$, show that $\tan\alpha = 2 + \sqrt{6} - \sqrt{3} - \sqrt{2}$. **[5]**

11 (a) Use the substitution $u^2 = x^2 + 3$ to show that $\int \frac{4x^3}{\sqrt{x^2+3}}\mathrm{d}x = \frac{4}{3}(x^2-6)\sqrt{x^2+3} + c$. **[5]**

(b) In this question you must show detailed reasoning.



The graph shows part of the curve $y = \frac{4x^3}{\sqrt{x^2+2}}$.

Find the exact area enclosed by the curve $y = \frac{4x^3}{\sqrt{x^2+3}}$, the normal to this curve at the point (1, 2) and the x-axis. **[7]**

stemup.co.uk

12 A cake is cooling so that, t minutes after it is removed from an oven, its temperature is θ °C. When the cake is removed from the oven, its temperature is 160 °C. After 10 minutes its temperature has fallen to 125 °C.

(a) In a simple model, the rate of decrease of the temperature of the cake is assumed to be constant.

(i) Write down a differential equation for this model. **[1]**

(ii) Solve this differential equation to find θ in terms of t. **[2]**

(iii) State **one** limitation of this model. **[1]**

(b) In a revised model, the rate of decrease of the temperature of the cake is proportional to the difference between the temperature of the cake and the temperature of the room. The temperature of the room is a constant 20 °C.

(i) Write down a differential equation for this revised model. **[1]**

(ii) Solve this differential equation to find θ in terms of t. **[6]**

(c) The cake can be decorated when its temperature is 25 °C. Find the difference in time between when the two models would predict that the cake can be decorated, giving your answer correct to the nearest minute. **[2]**

END OF QUESTION PAPER

OCR
Oxford Cambridge and RSA

Copyright Information

OCR is committed to seeking permission to reproduce all third-party content that it uses in its assessment materials. OCR has attempted to identify and contact all copyright holders whose work is used in this paper. To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced in the OCR Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download from our public website (www.ocr.org.uk) after the live examination series.

If OCR has unwittingly failed to correctly acknowledge or clear any third-party content in this assessment material, OCR will be happy to correct its mistake at the earliest possible opportunity.

For queries or further information please contact The OCR Copyright Team, The Triangle Building, Shaftesbury Road, Cambridge CB2 8EA.

OCR is part of the Cambridge Assessment Group; Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.

© OCR 2021

stemup.co.uk

Surname	StemUp Model Solution
Answer booklet	Write the question number in the left-hand margin at the start of each answer.

Question number

1

Two distinct real roots $\Rightarrow$ $b^2 - 4ac > 0$

$4^2 - 4 \times 1 \times (k + 3) > 0$

$16 - 4k - 12 > 0$

$4 - 4k > 0 \quad \Rightarrow \quad 4k < 4$

$k < 1$

2 (a)

Working in pence ($m > 100$):

Contract A: $C = 4000 + 4m$

Contract B: $C = 6(m - 100)$

2 (b)

$4000 + 4m = 6(m - 100)$

$4000 + 4m = 6m - 600$

$2m = 4600$

$m = 2300$ (minutes)

3 (a)

$x = ky^2\sqrt{z}$

$30 = k \times 2^2 \times \sqrt{9} = 12k \quad \Rightarrow \quad k = 2.5$

$x = 2.5y^2\sqrt{z}$

3 (b)

$x = 2.5 \times 3^2 \times \sqrt{25} = 2.5 \times 9 \times 5$

$x = 112.5$

4 (a)

$f(0.5) = 2(0.5)^3 - 3(0.5)^2 - 11(0.5) + 6$

$= 0.25 - 0.75 - 5.5 + 6 = 0$

$f(\frac{1}{2}) = 0$, so by the factor theorem $(2x - 1)$ is a factor of $f(x)$.

Question number	
4 (b)	$2x^3 - 3x^2 - 11x + 6 = (2x - 1)(x^2 + bx + c)$
	Comparing x^2: $2b - 1 = -3 \Rightarrow b = -1$; constants: $-c = 6 \Rightarrow c = -6$
	Check x: $-b + 2c = 1 - 12 = -11$ ✓
	$f(x) = (2x - 1)(x^2 - x - 6)$
	$f(x) = (2x - 1)(x - 3)(x + 2)$
4 (c)	$8^y = (2^y)^3$ and $4^y = (2^y)^2$, so let $x = 2^y$:
	$2x^3 - 3x^2 - 11x + 6 = 0 \Rightarrow (2x - 1)(x - 3)(x + 2) = 0$
	$x = 0.5$, $x = 3$ or $x = -2$
	$2^y = 0.5 \Rightarrow y = -1$
	$2^y = 3 \Rightarrow y = \log_2 3 = \frac{\ln 3}{\ln 2} = 1.58$ (3 s.f.)
	$2^y = -2$ has no solutions, since $2^y > 0$ for all y
	Hence $y = -1$ or $y = \log_2 3$
5(a)(i)	(2, 9)
5(a)(ii)	$3x - 1 = 2 \Rightarrow x = 1$; $y = 2 \times 6 = 12$
	(1, 12)
5(a)(iii)	(6, 2)
5(b)(i)	Stationary points where $g'(x) = 0$: $x = -2$ and $x = 4$
5(b)(ii)	$g(x)$ decreasing where $g'(x) < 0$: $x < -2$
5(b)(iii)	At $x = 4$, $g'(x)$ has a minimum (gradient of g' is zero) but g' does
	not change sign, so it is a stationary point of inflection: $x = 4$

Question number

6 (a)

$(8 - 3x)^{1/3} = 8^{1/3}(1 - \frac{3}{8}x)^{1/3} = 2(1 - \frac{3}{8}x)^{1/3}$

$(1 - \frac{3}{8}x)^{1/3} = 1 + (\frac{1}{3})(-\frac{3}{8}x) + \frac{(1/3)(-2/3)}{2!}(-\frac{3}{8}x)^2 + \ldots$

$= 1 - \frac{1}{8}x - \frac{1}{9} \times \frac{9}{64}x^2 + \ldots = 1 - \frac{1}{8}x - \frac{1}{64}x^2 + \ldots$

$(8 - 3x)^{1/3} = 2 - \frac{1}{4}x - \frac{1}{32}x^2 + \ldots$

6 (b)

Valid for $|\frac{3}{8}x| < 1$, i.e. $|x| < \frac{8}{3}$

6 (c)

$(1 + 2x)^{-2} = 1 + (-2)(2x) + \frac{(-2)(-3)}{2!}(2x)^2 + \ldots = 1 - 4x + 12x^2 + \ldots$

$(2 - \frac{1}{4}x - \frac{1}{32}x^2)(1 - 4x + 12x^2 + \ldots)$

Coefficient of x^2: $2 \times 12 + (-\frac{1}{4}) \times (-4) + (-\frac{1}{32}) \times 1$

$= 24 + 1 - \frac{1}{32} = \frac{799}{32}$ $(= 24.96875)$

7 (a)

$y = (x^2 - 2)\ln x \Rightarrow \frac{dy}{dx} = 2x \ln x + (x^2 - 2) \times \frac{1}{x} = 2x \ln x + \frac{x^2 - 2}{x}$

At a stationary point $\frac{dy}{dx} = 0$: $2x \ln x + \frac{x^2 - 2}{x} = 0$

Multiply by x: $2x^2 \ln x + x^2 - 2 = 0$ as required

Question number

7 (b) Let $f(x) = 2x^2 \ln x + x^2 - 2$

$f'(x) = 4x \ln x + 2x^2 \times \frac{1}{x} + 2x = 4x \ln x + 4x$

$x_{n+1} = x_n - \frac{2x_n^2 \ln x_n + x_n^2 - 2}{4x_n \ln x_n + 4x_n}$

$x_{n+1} = \frac{x_n(4x_n \ln x_n + 4x_n) - (2x_n^2 \ln x_n + x_n^2 - 2)}{4x_n \ln x_n + 4x_n}$

$x_{n+1} = \frac{4x_n^2 \ln x_n + 4x_n^2 - 2x_n^2 \ln x_n - x_n^2 + 2}{4x_n \ln x_n + 4x_n}$

$x_{n+1} = \frac{2x_n^2 \ln x_n + 3x_n^2 + 2}{4x_n(\ln x_n + 1)}$ as required

7 (c) $x_2 = \frac{0 + 3 + 2}{4(0 + 1)} = 1.25$

$x_3 = 1.207515437... = 1.2075$ (4 d.p.)

7 (d) $x_4 = 1.206124696...$, $x_5 = 1.206123218...$, $x_6 = 1.206123218...$

$x = 1.206$ (3 d.p.)

$y = (1.206123...^2 - 2) \ln 1.206123... = -0.102189... = -0.102$

Stationary point $(1.206, -0.102)$

8(a)(i) $f(x) \in \mathbb{R}$ (all real values)

8(a)(ii) $g(x) \leq -1$ or $g(x) \geq 1$, i.e. $g(x) \in (-\infty, -1] \cup [1, \infty)$

8(b)(i) $\cos 0.6 = 0.825335...$, so $g(0.6) = \sec 0.6 = 1 \div 0.825335... = 1.211628...$

$fg(0.6) = 2 \tan(1.211628...) = 2 \times 2.663447... = 5.326894...$

$fg(0.6) = 5.33$ (3 s.f.) as required

Question number

8(b)(ii) f is a many-to-one function on $0 \leqslant x \leqslant 2\pi$ (e.g. $\tan x$ repeats every π),

so f has no inverse and f^{-1} is not defined.

8 (c) $4\tan^2 x + 6\sec x = 0$

$4(\sec^2 x - 1) + 6\sec x = 0$

$4\sec^2 x + 6\sec x - 4 = 0 \quad \Rightarrow \quad 2\sec^2 x + 3\sec x - 2 = 0$

$(2\sec x - 1)(\sec x + 2) = 0$

$\sec x = \frac{1}{2} \quad$ or $\quad \sec x = -2$

$\sec x = \frac{1}{2}$ has no solutions, since $|\sec x| \geqslant 1$

$\sec x = -2 \Rightarrow \cos x = -\frac{1}{2} \Rightarrow x = \frac{2\pi}{3}, \frac{4\pi}{3}$

9 (a) $e^{-3t} > 0$ for all t, so $y = 2e^{-3t} > 0$ for all $t \geqslant 0$.

y is never 0, so the path never crosses the x-axis.

9 (b) On the y-axis $x = 0$: $e^{2t} - 4e^t + 3 = 0$

$(e^t - 1)(e^t - 3) = 0 \quad \Rightarrow \quad e^t = 1$ or $e^t = 3$

$t = 0$ or $t = \ln 3$

9 (c) $\frac{dx}{dt} = 2e^{2t} - 4e^t, \qquad \frac{dy}{dt} = -6e^{-3t}$

$\frac{dy}{dx} = \frac{dy/dt}{dx/dt} = \frac{-6e^{-3t}}{2e^{2t} - 4e^t} = \frac{-3e^{-3t}}{e^{2t} - 2e^t}$

$= \frac{-3}{e^{3t}(e^{2t} - 2e^t)} = \frac{-3}{e^{5t} - 2e^{4t}} = \frac{3}{2e^{4t} - e^{5t}}$ as required

Question number

9 (d)

Parallel to the y-axis when $\frac{dx}{dy} = 0$, i.e. $2e^{4t} - e^{5t} = 0$

$e^{4t}(2 - e^{t}) = 0$; $e^{4t} \neq 0$ so $e^{t} = 2$, $t = \ln 2$

$x = e^{2\ln 2} - 4e^{\ln 2} + 3 = 4 - 8 + 3 = -1$

$y = 2e^{-3\ln 2} = 2 \times \frac{1}{8} = \frac{1}{4}$ Coordinates $(-1, \frac{1}{4})$

10(a)(i)

Triangle BCD is right-angled at D, so $\cos y = \frac{CD}{BC} = \frac{CD}{a}$

Hence $CD = a \cos y$

10(a)(ii)

Area ADC $= \frac{1}{2} \times AC \times CD \times \sin(\text{angle } ACD)$

$= \frac{1}{2} \times b \times (a \cos y) \times \sin x = \frac{1}{2} ab \sin x \cos y$ as required

10(a)(iii)

Similarly, from triangle ADC, $\cos x = CD/b$, so $CD = b \cos x$

Area BCD $= \frac{1}{2} \times BC \times CD \times \sin y = \frac{1}{2} a(b \cos x) \sin y = \frac{1}{2} ab \cos x \sin y$

Area ABC $= \frac{1}{2} \times AC \times BC \times \sin(\text{angle } ACB) = \frac{1}{2} ab \sin(x + y)$

Area ABC = area ADC + area BCD:

$\frac{1}{2} ab \sin(x + y) = \frac{1}{2} ab \sin x \cos y + \frac{1}{2} ab \cos x \sin y$

Divide by ½ab: $\sin(x + y) = \sin x \cos y + \cos x \sin y$ as required

Question number

10 (b) $\sin 30° \cos \alpha + \cos 30° \sin \alpha = \cos 45° \cos \alpha + \sin 45° \sin \alpha$

$\frac{1}{2} \cos \alpha + \frac{\sqrt{3}}{2} \sin \alpha = \frac{\sqrt{2}}{2} \cos \alpha + \frac{\sqrt{2}}{2} \sin \alpha$

Multiply by 2: $\cos \alpha + \sqrt{3} \sin \alpha = \sqrt{2} \cos \alpha + \sqrt{2} \sin \alpha$

$(\sqrt{3} - \sqrt{2}) \sin \alpha = (\sqrt{2} - 1) \cos \alpha$

$\frac{\sin \alpha}{\cos \alpha} = \tan \alpha = \frac{\sqrt{2} - 1}{\sqrt{3} - \sqrt{2}} = \frac{(\sqrt{2} - 1)(\sqrt{3} + \sqrt{2})}{(\sqrt{3} - \sqrt{2})(\sqrt{3} + \sqrt{2})}$

$\tan \alpha = \frac{\sqrt{6} + 2 - \sqrt{3} - \sqrt{2}}{3 - 2} = 2 + \sqrt{6} - \sqrt{3} - \sqrt{2}$ as required

11 (a) $u^2 = x^2 + 3 \Rightarrow 2u\,du = 2x\,dx \Rightarrow x\,dx = u\,du$

$\int \frac{4x^3}{\sqrt{(x^2 + 3)}}\,dx = \int \frac{4x^2 \times x}{\sqrt{(x^2 + 3)}}\,dx = \int \frac{4(u^2 - 3)}{u} \times u\,du$

$= \int (4u^2 - 12)\,du$

$= \frac{4}{3} u^3 - 12u + c = \frac{4}{3} u(u^2 - 9) + c$

$= \frac{4}{3} (x^2 + 3 - 9)\sqrt{(x^2 + 3)} + c = \frac{4}{3} (x^2 - 6)\sqrt{(x^2 + 3)} + c$ as required

Question number

11 (b) Area under curve from $x = 0$ to $x = 1$ ($u = \sqrt{3}$ to $u = 2$):

$$\int_0^1 \frac{4x^3}{\sqrt{(x^2+3)}}\,dx = \left[\frac{4}{3}(x^2-6)\sqrt{(x^2+3)}\right]_0^1 = \frac{4}{3}((-5)(2) - (-6)(\sqrt{3}))$$

$$= \frac{4}{3}(6\sqrt{3} - 10) = 8\sqrt{3} - \frac{40}{3} \quad (= 0.523\ldots)$$

$$\frac{dy}{dx} = \frac{12x^2(x^2+3)^{1/2} - 4x^3 \times x(x^2+3)^{-1/2}}{x^2+3}$$

At $x = 1$: $\frac{dy}{dx} = \frac{12 \times 2 - 4 \times \frac{1}{2}}{4} = \frac{22}{4} = \frac{11}{2}$, so gradient of normal $= -\frac{2}{11}$

Normal: $y - 2 = -\frac{2}{11}(x-1)$; $y = 0$: $-22 = -2(x-1) \Rightarrow x = 12$

Triangle under normal: $\frac{1}{2} \times (12-1) \times 2 = 11$

Total area $= 8\sqrt{3} - \frac{40}{3} + 11 = 8\sqrt{3} - \frac{7}{3}$

12(a)(i) $\frac{d\theta}{dt} = -k$, where k is a positive constant

12(a)(ii) $\theta = -kt + c$; $t = 0$, $\theta = 160 \Rightarrow c = 160$

$t = 10$, $\theta = 125$: $125 = 160 - 10k \Rightarrow k = 3.5$

$\theta = 160 - 3.5t$

12(a)(iii) The model predicts the temperature keeps falling forever, below room temperature and eventually below freezing point, which is unrealistic.

12(b)(i) $\frac{d\theta}{dt} = -k(\theta - 20)$, where k is a positive constant

Question number

12(b)(ii)

$\int \frac{1}{\theta - 20}\, d\theta = \int -k\, dt$

$\ln|\theta - 20| = -kt + c$

$t = 0,\ \theta = 160$: $\ln 140 = c$

$t = 10,\ \theta = 125$: $\ln 105 = -10k + \ln 140 \Rightarrow k = -0.1 \ln \frac{105}{140} = -0.1 \ln 0.75$

$(k = 0.028768\ldots)$

$\ln(\theta - 20) = (0.1 \ln 0.75)t + \ln 140$

$\theta - 20 = e^{(0.1 \ln 0.75)t + \ln 140} = 140e^{(0.1 \ln 0.75)t}$

$\theta = 20 + 140e^{(0.1 \ln 0.75)t}$ (or $\theta = 20 + 140(0.75)^{0.1t}$)

12 (c)

Simple model: $25 = 160 - 3.5t \Rightarrow t = 38.571\ldots$ minutes

Revised model: $5 = 140e^{(0.1 \ln 0.75)t} \Rightarrow \ln 5 = (0.1 \ln 0.75)t + \ln 140$

$t = \frac{\ln(5/140)}{0.1 \ln 0.75} = 115.829\ldots$ minutes

Difference $= 115.829\ldots - 38.571\ldots = 77.26\ldots \approx 77$ minutes