OCR
Oxford Cambridge and RSA

Wednesday 15 May 2019 – Morning

AS Level Mathematics A

H230/01 Pure Mathematics and Statistics

Time allowed: 1 hour 30 minutes

7698691988

You must have:
- Printed Answer Booklet

You may use:
- a scientific or graphical calculator

INSTRUCTIONS
- Use black ink. HB pencil may be used for graphs and diagrams only.
- Answer **all** the questions.
- **Write your answer to each question in the space provided in the Printed Answer Booklet.** If additional space is required, you should use the lined page(s) at the end of the Printed Answer Booklet. The question number(s) must be clearly shown.
- You are permitted to use a scientific or graphical calculator in this paper.
- Give non-exact numerical answers correct to 3 significant figures unless a different degree of accuracy is specified in the question.
- The acceleration due to gravity is denoted by $g\,\mathrm{m\,s^{-2}}$. Unless otherwise instructed, when a numerical value is needed, use $g = 9.8$.

INFORMATION
- The total mark for this paper is **75**.
- The marks for each question are shown in brackets **[ ]**.
- **You are reminded of the need for clear presentation in your answers.**
- The Printed Answer Booklet consists of **16** pages. The Question Paper consists of **8** pages.

© OCR 2019 [603/0933/7]
DC (KN/FC) 171474/3

OCR is an exempt Charity

stemup.co.uk

Formulae
AS Level Mathematics A (H230)

Binomial series

$$(a+b)^n = a^n + {}^n\mathrm{C}_1 a^{n-1}b + {}^n\mathrm{C}_2 a^{n-2}b^2 + \ldots + {}^n\mathrm{C}_r a^{n-r}b^r + \ldots + b^n \qquad (n \in \mathbb{N}),$$

where ${}^n\mathrm{C}_r = {}_n\mathrm{C}_r = \binom{n}{r} = \dfrac{n!}{r!(n-r)!}$

Differentiation from first principles

$$\mathrm{f}'(x) = \lim_{h \to 0} \frac{\mathrm{f}(x+h) - \mathrm{f}(x)}{h}$$

Standard deviation

$$\sqrt{\frac{\Sigma(x-\overline{x})^2}{n}} = \sqrt{\frac{\Sigma x^2}{n} - \overline{x}^2} \quad \text{or} \quad \sqrt{\frac{\Sigma f(x-\overline{x})^2}{\Sigma f}} = \sqrt{\frac{\Sigma f x^2}{\Sigma f} - \overline{x}^2}$$

The binomial distribution

If $X \sim \mathrm{B}(n, p)$ then $P(X = x) = \binom{n}{x} p^x (1-p)^{n-x}$, mean of X is np, variance of X is $np(1-p)$

Kinematics

$$v = u + at$$

$$s = ut + \tfrac{1}{2}at^2$$

$$s = \tfrac{1}{2}(u+v)t$$

$$v^2 = u^2 + 2as$$

$$s = vt - \tfrac{1}{2}at^2$$

© OCR 2019

stemup.co.uk

Section A: Pure Mathematics

Answer **all** the questions.

1 It is given that $\mathrm{f}(x) = 3x - \frac{5}{x^3}$.

Find

(a) $\mathrm{f}'(x)$, **[3]**

(b) $\mathrm{f}''(x)$, **[2]**

(c) $\int \mathrm{f}(x)\,\mathrm{d}x$. **[3]**

2 The circle $x^2 + y^2 - 4x + ky + 12 = 0$ has radius 1.

Find the two possible values of the constant k. **[4]**

3 **In this question you must show detailed reasoning.**

(a) The polynomial $\mathrm{f}(x)$ is defined by $\mathrm{f}(x) = 2x^3 + 3x^2 - 8x + 3$.

(i) Show that $\mathrm{f}(1) = 0$. **[1]**

(ii) Solve the equation $\mathrm{f}(x) = 0$. **[4]**

(b) Hence solve the equation $2\sin^3\theta + 3\sin^2\theta - 8\sin\theta + 3 = 0$ for $0° \leqslant \theta < 360°$. **[5]**

4 **(a)** Find the coordinates of the stationary points on the curve $y = x^3 - 6x^2 + 9x$. **[4]**

(b) The equation $x^3 - 6x^2 + 9x + k = 0$ has exactly one real root.

Using your answers from part **(a)** or otherwise, find the range of possible values of k. **[2]**

stemup.co.uk


5 **(a)** Prove that the following statement is **not** true.

m is an odd number greater than $1 \Rightarrow m^2 + 4$ is prime. **[1]**

(b) By considering separately the case when n is odd and the case when n is even, prove that the following statement **is** true.

n is a positive integer $\Rightarrow n^2 + 1$ is not a multiple of 4. **[4]**

6



The diagram shows triangle ABC, with $AB = x$ cm, $AC = y$ cm and angle $BAC = 60°$. It is given that the area of the triangle is $(x+y)\sqrt{3}$ cm^2.

(a) Show that $4x + 4y = xy$. **[2]**

When the vertices of the triangle are placed on the circumference of a circle, AC is a diameter of the circle.

(b) Determine the value of x and the value of y. **[4]**

7 **(a)** Write down an expression for the gradient of the curve $y = e^{kx}$. **[1]**

(b) The line L is a tangent to the curve $y = e^{\frac{1}{2}x}$ at the point where $x = 2$.

Show that L passes through the point (0, 0). **[4]**

(c) Find the coordinates of the point of intersection of the curves $y = 3e^x$ and $y = 1 - 2e^{\frac{1}{2}x}$. **[6]**

© OCR 2019

stemup.co.uk

Section B: Statistics

Answer **all** the questions.

8 **(a)** Joseph drew a histogram to show information about one Local Authority. He used data from the "Age structure by LA 2011" tab in the large data set. The table shows an extract from the data that he used.

Age group	0 to 4
Frequency	2143

Joseph used a scale of 1 cm = 1000 units on the frequency density axis.

Calculate the height of the histogram block for the 0 to 4 class. **[2]**

(b) Magdalene wishes to draw a statistical diagram to illustrate some of the data from the "Method of travel by LA 2011" tab in the large data set.

State why she cannot draw a histogram. **[1]**

9 The table shows information about the number of days absent last year by students in class 2A at a certain school.

Number of days absent	0	1	2 to 4	5 to 10	11 to 20	21 to 30	More than 30
Number of students	7	12	9	1	0	1	0

(a) Calculate an estimate of the mean for these data. **[2]**

(b) Find the median of these data. **[1]**

The headteacher is writing a report on the numbers of absences at her school. She wishes to include a figure for the average number of absences in class 2A. A governor suggests that she should quote the mean. The class teacher suggests that she should quote the median, because it is lower than the mean.

(c) Give another reason for using the median rather than the mean for the average number of absences in class 2A. **[1]**

stemup.co.uk

10 The table shows extracts from the "Method of travel by LA" tabs for 2001 and 2011 in the large data set.

Local authority (LA)	All people in employment	Underground, metro, light rail, tram	Train	Bus, minibus or coach	Motorcycle, scooter or moped	Driving a car or van
LA1 2001	79 226	14 369	5235	20 575	1227	16 052
LA1 2011	118 556	22 486	8336	30 541	1220	12 445
LA2 2001	203 614	190	1062	15 327	1256	121 690
LA2 2011	227 894	323	1865	13 732	1038	146 644
LA3 2001	42 993	35	482	4363	274	24 105
LA3 2011	49 014	33	828	3380	191	28 981
LA4 2001	101 697	65	693	21 758	846	45 407
LA4 2011	123 218	2495	1315	24 275	763	54 020

(a) In one of these four LAs a new tram system was opened in 2004.

Suggest, with a reason taken from the data, which LA this could have been. **[2]**

(b) Julian suggests that the figures for "Bus, minibus or coach" for LA1 show that some new bus routes were probably introduced in this LA between 2001 and 2011.

Use data from the table to comment on this suggestion. **[2]**

(c) In one of these four LAs a congestion charge on vehicles was introduced in 2003.

Suggest, with a reason taken from the data, which LA this could have been. **[2]**

11 It is known that, under the standard treatment for a certain disease, 9.7% of patients with the disease experience side effects within one year.

In a trial of a new treatment, a random sample of 450 patients with this disease was selected and the number X who experienced side effects within one year was noted.

(a) State one assumption needed in order to use a binomial model for X. **[1]**


It was found that 51 of the 450 patients experienced side effects within one year.

(b) Test, at the 10% significance level, whether the proportion of patients experiencing side effects within one year is greater under the new treatment than under the standard treatment. **[7]**

© OCR 2019

stemup.co.uk

12 The Venn diagram shows the numbers of students studying various subjects, in a year group of 100 students.



A student is chosen at random from the 100 students. Then another student is chosen from the remaining students.

Find the probability that the first student studies History and the second student studies Geography but not Psychology. **[4]**

END OF QUESTION PAPER

© OCR 2019

stemup.co.uk

OCR
Oxford Cambridge and RSA

Copyright Information

OCR is committed to seeking permission to reproduce all third-party content that it uses in its assessment materials. OCR has attempted to identify and contact all copyright holders whose work is used in this paper. To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced in the OCR Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download from our public website (www.ocr.org.uk) after the live examination series.

If OCR has unwittingly failed to correctly acknowledge or clear any third-party content in this assessment material, OCR will be happy to correct its mistake at the earliest possible opportunity.

For queries or further information please contact The OCR Copyright Team, The Triangle Building, Shaftesbury Road, Cambridge CB2 8EA.

OCR is part of the Cambridge Assessment Group; Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.

stemup.co.uk

Surname	StemUp Model Solution
Answer booklet	Write the question number in the left-hand margin at the start of each answer.

Question number

1 (a)

$f(x) = 3x - 5x^{-3}$

$f'(x) = 3 + 15x^{-4} = 3 + \frac{15}{x^4}$

1 (b)

$f''(x) = -60x^{-5} = -\frac{60}{x^5}$

1 (c)

$\int f(x)\,dx = \int (3x - 5x^{-3})\,dx$

$= \frac{3x^2}{2} - \frac{5x^{-2}}{-2} + c$

$= \frac{3x^2}{2} + \frac{5}{2x^2} + c$

2

$x^2 - 4x + y^2 + ky + 12 = 0$

$(x-2)^2 - 4 + \left(y + \frac{k}{2}\right)^2 - \frac{k^2}{4} + 12 = 0$

$(x-2)^2 + \left(y + \frac{k}{2}\right)^2 = \frac{k^2}{4} - 8$

Radius 1: $\frac{k^2}{4} - 8 = 1 \Rightarrow \frac{k^2}{4} = 9$

$k^2 = 36$

$k = 6$ or $k = -6$

3 (a)(i)

$f(1) = 2(1)^3 + 3(1)^2 - 8(1) + 3 = 2 + 3 - 8 + 3 = 0$

3 (a)(ii)

$f(1) = 0$ so $(x - 1)$ is a factor of $f(x)$

$2x^3 + 3x^2 - 8x + 3 = (x-1)(2x^2 + 5x - 3)$

$= (x-1)(2x-1)(x+3)$

$f(x) = 0 \Rightarrow x = 1,\ x = \frac{1}{2},\ x = -3$

3 (b)

Let $x = \sin\theta$: $2x^3 + 3x^2 - 8x + 3 = 0$

so $\sin\theta = 1$, $\sin\theta = \frac{1}{2}$ or $\sin\theta = -3$

Question number	
	$\sin\theta = 1 \Rightarrow \theta = 90°$
	$\sin\theta = \frac{1}{2} \Rightarrow \theta = 30°$ or $\theta = 180° - 30° = 150°$
	$\sin\theta = -3$ has no solutions since $-1 \leq \sin\theta \leq 1$
	$\theta = 30°, 90°, 150°$
4 (a)	$y = x^3 - 6x^2 + 9x \Rightarrow \frac{dy}{dx} = 3x^2 - 12x + 9$
	Stationary points: $3x^2 - 12x + 9 = 0$
	$x^2 - 4x + 3 = 0 \Rightarrow (x-1)(x-3) = 0 \Rightarrow x = 1$ or $x = 3$
	$x = 1$: $y = 1 - 6 + 9 = 4$ $x = 3$: $y = 27 - 54 + 27 = 0$
	Stationary points are $(1, 4)$ and $(3, 0)$
4 (b)	$x^3 - 6x^2 + 9x = -k$: the line $y = -k$ meets the curve once
	only if it lies above the max $(1, 4)$ or below the min $(3, 0)$
	so $-k > 4$ or $-k < 0$
	$k < -4$ or $k > 0$
5 (a)	Counter-example: $m = 9$ is odd and greater than 1, but
	$m^2 + 4 = 81 + 4 = 85 = 5 \times 17$, which is not prime.
	So the statement is not true.
5 (b)	n even: $n = 2k$ (k a positive integer)
	$n^2 + 1 = 4k^2 + 1$, which is 1 more than a multiple of 4,
	so it is not a multiple of 4.
	n odd: $n = 2k + 1$ (k an integer, $k \geq 0$)
	$n^2 + 1 = 4k^2 + 4k + 1 + 1 = 4(k^2 + k) + 2$
	$4(k^2 + k)$ is a multiple of 4, so $4(k^2 + k) + 2$ leaves remainder 2
	when divided by 4, so it is not a multiple of 4.
	Every positive integer is odd or even, $\therefore n^2 + 1$ is never a multiple of 4.

Question number	
6 (a)	Area $= \frac{1}{2}xy\sin 60° = \frac{1}{2} \times \frac{\sqrt{3}}{2} \times xy = \frac{\sqrt{3}}{4}xy$
	$\frac{\sqrt{3}}{4}xy = (x + y)\sqrt{3} \quad \Rightarrow \quad xy = 4(x + y)$
	$\Rightarrow \quad 4x + 4y = xy$ as required
6 (b)	AC is a diameter, so angle ABC = 90° (angle in a semicircle)
	$\cos 60° = \frac{x}{y} = \frac{1}{2} \quad \Rightarrow \quad y = 2x$
	Substitute into $4x + 4y = xy$: $4x + 8x = 2x^2$
	$2x^2 - 12x = 0 \quad \Rightarrow \quad 2x(x - 6) = 0 \quad \Rightarrow \quad x = 6 \quad (x \neq 0)$
	$x = 6, \quad y = 12$
7 (a)	$\frac{dy}{dx} = ke^{kx}$
7 (b)	$y = e^{x/2} \quad \Rightarrow \quad \frac{dy}{dx} = \frac{1}{2}e^{x/2}$
	At $x = 2$: gradient $= \frac{1}{2}e^1 = \frac{1}{2}e$, $\quad y = e^1 = e$
	Tangent: $y - e = \frac{1}{2}e(x - 2)$
	$y = \frac{1}{2}ex - e + e \quad \Rightarrow \quad y = \frac{1}{2}ex$
	When $x = 0$, $y = 0$, so L passes through (0, 0)
7 (c)	$3e^x = 1 - 2e^{x/2}$
	Let $u = e^{x/2}$, so $e^x = u^2$: $\quad 3u^2 + 2u - 1 = 0$
	$(3u - 1)(u + 1) = 0 \quad \Rightarrow \quad u = \frac{1}{3}$ or $u = -1$
	$e^{x/2} = -1$ has no solutions, since $e^{x/2} > 0$ for all x
	$e^{x/2} = \frac{1}{3} \quad \Rightarrow \quad \frac{x}{2} = \ln\frac{1}{3} \quad \Rightarrow \quad x = 2\ln\frac{1}{3} = \ln\frac{1}{9}$
	$y = 3e^x = 3 \times \frac{1}{9} = \frac{1}{3}$
	Point of intersection is $\left(\ln\frac{1}{9}, \frac{1}{3}\right)$ $\quad (= (-2\ln 3, \frac{1}{3}))$
8 (a)	Class 0 to 4 has width 5 (ages 0 up to 5)
	Frequency density $= \frac{2143}{5} = 428.6$
	Height $= \frac{428.6}{1000} = 0.4286 = 0.429$ cm (3 s.f.)

Question number	
8 (b)	The data are non-numerical (qualitative): the methods of travel are categories, not numerical values that can be grouped into classes.
9 (a)	Midpoints: 0, 1, 3, 7.5, 15.5, 25.5 $\Sigma fx = 0 + 12 + 27 + 7.5 + 0 + 25.5 = 72, \quad \Sigma f = 30$ Mean $= \frac{\Sigma fx}{\Sigma f} = \frac{72}{30} = 2.4$ days
9 (b)	$n = 30$: median is between the 15th and 16th values, both 1 Median = 1
9 (c)	The median is not affected by the outlier (the one student absent for 21 to 30 days), whereas this one value increases the mean.
10 (a)	LA4: the number travelling by underground, metro, light rail, tram increased from 65 to 2495, a far larger increase than in the other LAs.
10 (b)	Bus in LA1 rose from 20 575 to 30 541, but the total in employment also rose, from 79 226 to 118 556. Proportion by bus: $\frac{20\,575}{79\,226} = 0.260$ and $\frac{30\,541}{118\,556} = 0.258$ The proportion is about the same, so the rise is due to the increase in total workers; the data give no reason to suggest new bus routes.
10 (c)	LA1: the number driving a car or van decreased (16 052 to 12 445) despite the number in employment increasing (79 226 to 118 556); in all the other LAs the number driving increased.
11 (a)	Patients experience side effects independently of each other.
11 (b)	Let p = proportion of patients (under the new treatment) who

Question number

experience side effects within one year.

H_0: $p = 0.097$ H_1: $p > 0.097$

Under H_0, $X \sim B(450, 0.097)$

$P(X \geqslant 51) = 1 - P(X \leqslant 50) = 1 - 0.86170...$

$= 0.13830... = 0.138$ (3 s.f.)

$0.138 > 0.1$, so do not reject H_0

There is insufficient evidence at the 10% level that the proportion of patients experiencing side effects within one year is greater under the new treatment than under the standard treatment.

12

$n(H) = 7 + 25 + 3 + 8 = 43$, $n(G \cap P') = 25 + 12 = 37$

First in $H \cap G \cap P'$ (25): second from 36 remaining in $G \cap P'$

First in $H \cap G \cap P$ (3) or $H \cap G'$ $(7 + 8 = 15)$: second from 37

$$P = \frac{25}{100} \times \frac{36}{99} + \frac{3}{100} \times \frac{37}{99} + \frac{15}{100} \times \frac{37}{99}$$

$$= \frac{900 + 111 + 555}{9900} = \frac{1566}{9900}$$

$$= \frac{87}{550} = 0.158 \text{ (3 s.f.)}$$