

Please write clearly in block capitals.

Centre number

--	--	--	--	--

Candidate number

--	--	--	--

Surname *StemUp Model Solution*

Forename(s)

Candidate signature

I declare this is my own work.

INTERNATIONAL A-LEVEL CHEMISTRY (9620)

Unit 3: Inorganic 2 and Physical 2

Thursday 12 January 2023 07:00 GMT Time allowed: 1 hour 30 minutes

Materials

For this paper you must have:

- the Periodic Table/Data Sheet, provided as an insert
- a ruler with millimetre measurements
- a scientific calculator, which you are expected to use where appropriate.

Instructions

- Use black ink or black ball-point pen.
- Fill in the boxes at the top of this page.
- Answer **all** questions.
- You must answer the questions in the spaces provided. Do not write outside the box around each page or on blank pages.
- If you need extra space for your answer(s), use the lined pages at the end of this book. Write the question number against your answer(s).
- All working must be shown.
- Do all rough work in this book. Cross through any work you do not want to be marked.

Information

- The marks for questions are shown in brackets.
- The maximum mark for this paper is 80.

For Examiner's Use	
Question	Mark
1	
2	
3	
4	
5	
6	
7	
8	
TOTAL	



J A N 2 3 C H 0 3 0 1

Answer **all** questions in the spaces provided.

0 1

This question is about the enthalpy of lattice formation of barium chloride.

0 1 . 1

Define enthalpy of lattice formation.

[2 marks]


The enthalpy change when 1 mole of a solid ionic compound (lattice) is formed from its gaseous ions (under standard conditions).

0 1 . 2

Figure 1 shows an incomplete Born–Haber cycle for barium chloride, BaCl_2

The Born–Haber cycle is not to scale.

Figure 1



Write the formulas, including state symbols, of the appropriate species on each of the two blank horizontal lines in **Figure 1**.

[2 marks]



0 1 . 3

Table 1 shows enthalpy changes from the Born–Haber cycle for barium chloride.

Table 1

Enthalpy change	$\Delta H^\ominus / \text{kJ mol}^{-1}$
Enthalpy of atomisation of barium	+176
First ionisation enthalpy of barium	+502
Enthalpy of atomisation of chlorine	+121
First electron affinity of chlorine	–364
Enthalpy of lattice formation of barium chloride	–2018
Enthalpy of formation of barium chloride	–860

Use the data in **Table 1** to calculate the **second** ionisation enthalpy of barium.

Explain why this value for the **second** ionisation enthalpy of barium is different from the value for the **first** ionisation enthalpy of barium given in **Table 1**.

[4 marks]

Calculation

$$\Delta H_a(\text{Ba}) + 2\Delta H_a(\text{Cl}) + 1E_1 + 1E_2 + 2EA(\text{Cl}) + \Delta H_{LF} = \Delta H_f$$

$$176 + (2 \times 121) + 502 + 1E_2 + (2 \times -364) + (-2018) = -860$$

$$-1826 + 1E_2 = -860$$

$$1E_2 = -860 + 1826 = +966 \text{ kJ mol}^{-1}$$

Second ionisation enthalpy +966 kJ mol^{–1}

Explanation More energy is needed to remove an electron from a positive

Ba⁺ ion than from a neutral Ba atom.

Ba⁺ has more protons than electrons, so the remaining electrons are

held more strongly by the nucleus and are harder to remove.

Turn over ►



0 1 . 4

The enthalpy of lattice formation for barium chloride in **Table 1** was determined from experimental values.

The value calculated from theoretical data is almost the same as this experimental value.

Use your knowledge of the bonding in barium chloride to explain why these two values are almost the same.

[1 mark]

Barium chloride is (almost) perfectly ionic: the ions are spherical and Ba^{2+} does not polarise the Cl^- ions, so there is no covalent character.

0 1 . 5

Table 2 shows some enthalpy data for barium chloride and its ions.

Table 2

Enthalpy change	$\Delta H^\ominus / \text{kJ mol}^{-1}$
Enthalpy of hydration of $Ba^{2+}(g)$	-1360
Enthalpy of hydration of $Cl^-(g)$	-364
Enthalpy of lattice formation of barium chloride	-2018

Use the data in **Table 2** to calculate the standard enthalpy of solution of barium chloride.

[2 marks]

$$\Delta H_{\text{soln}} = -\Delta H_{\text{LF}} + \Delta H_{\text{hyd}}(Ba^{2+}) + 2\Delta H_{\text{hyd}}(Cl^-)$$

$$\Delta H_{\text{soln}} = -(-2018) + (-1360) + (2 \times -364)$$

$$\Delta H_{\text{soln}} = 2018 - 1360 - 728 = -70 \text{ kJ mol}^{-1}$$

Enthalpy of solution -70 kJ mol⁻¹

11



Turn over for the next question

*Do not write
outside the
box*

**DO NOT WRITE ON THIS PAGE
ANSWER IN THE SPACES PROVIDED**

Turn over ►



0 2

This question is about the thermodynamics of osmium (Os) and its oxide (OsO₄).

0 2 . 1

Which is the correct classification for the element osmium?

Tick (✓) **one** box.

[1 mark]

s block

☐

p block

☐

d block

☒

f block

☐

0 2 . 2

Osmium reacts slowly with atmospheric oxygen to form crystals of white osmium tetroxide, OsO₄




Table 3 shows thermodynamic data for this reaction.

Table 3

$\Delta G^\ominus / \text{kJ mol}^{-1}$	Temperature / K
−339	150
−250	450
−160	750
−71	1050
18	1350

Use these data to plot a graph of free-energy change, ΔG , (*y*-axis) against temperature (*x*-axis) on the grid in **Figure 2**.

Calculate the gradient of the line on your graph.


Use $\Delta G = \Delta H - T\Delta S$ and your gradient to calculate the entropy change (ΔS) in $\text{J K}^{-1} \text{mol}^{-1}$ for the reaction.

Determine the enthalpy change (ΔH) in kJ mol^{-1} for the reaction.

[5 marks]



Figure 2



$$\text{gradient} = (18 - (-339)) \div (1350 - 150) = 357 \div 1200 = 0.298 \text{ kJ K}^{-1} \text{ mol}^{-1}$$

$$\Delta G = -T\Delta S + \Delta H, \text{ so gradient} = -\Delta S: \Delta S = -0.298 \text{ kJ K}^{-1} \text{ mol}^{-1} = -298 \text{ J K}^{-1} \text{ mol}^{-1}$$

$$\Delta H = y\text{-intercept at } T = 0 \text{ (extrapolated line)} = -384 \text{ kJ mol}^{-1}$$

$$\text{check: } \Delta H = \Delta G + T\Delta S = 18 + 1350 \times (-0.298) = -384 \text{ kJ mol}^{-1}$$

$$\Delta S \quad \underline{-298} \quad \text{J K}^{-1} \text{ mol}^{-1}$$

$$\Delta H \quad \underline{-384} \quad \text{kJ mol}^{-1}$$

Turn over ►



0 2 . 3

Under different conditions, the same reactants form crystals of yellow osmium tetroxide.

The arrangement of the particles in crystals of yellow osmium tetroxide is different from the arrangement of the particles in crystals of white osmium tetroxide.

Table 4 shows the standard entropy values.

Table 4

Substance	$S^\ominus / \text{J K}^{-1} \text{mol}^{-1}$
yellow $\text{OsO}_4(\text{s})$	124
white $\text{OsO}_4(\text{s})$	145

Use the data in **Table 4** to explain the difference in the entropy of the two different forms of osmium tetroxide.

[1 mark]

Yellow OsO_4 has a lower entropy, so its particles are more ordered (less disordered) than in white OsO_4 .

0 2 . 4

The entropy change for the reaction between osmium and oxygen to form crystals of yellow osmium tetroxide is $-319 \text{ J K}^{-1} \text{mol}^{-1}$




Table 5 shows some standard entropy data.

Table 5

Substance	$S^\ominus / \text{J K}^{-1} \text{mol}^{-1}$
$\text{O}_2(\text{g})$	205
yellow $\text{OsO}_4(\text{s})$	124

Calculate the standard entropy, in $\text{J K}^{-1} \text{mol}^{-1}$, of the element osmium.

[2 marks]

$$\Delta S = S(\text{OsO}_4) - [S(\text{Os}) + 2S(\text{O}_2)]$$

$$-319 = 124 - S(\text{Os}) - (2 \times 205)$$


$$S(\text{Os}) = 124 - 410 + 319 = +33 \text{ J K}^{-1} \text{mol}^{-1}$$

$S^\ominus$ +33 $\text{J K}^{-1} \text{mol}^{-1}$



0 2 . 5

It is suggested that white osmium tetroxide can be reduced to osmium by reaction with hydrogen gas according to the equation



The entropy change for the reaction was calculated to be $-120 \text{ J K}^{-1} \text{ mol}^{-1}$

State whether this reaction is feasible at 298 K

Justify your answer with a calculation.

[2 marks]

$$\Delta G = \Delta H - T\Delta S = -584 - 298 \times (-120 \div 1000)$$

$$\Delta G = -584 + 35.8 = -548 \text{ kJ mol}^{-1}$$

The reaction is feasible at 298 K because ΔG is negative (less than zero).

11

Turn over for the next question

Turn over ►



0 3

This question is about acids, bases and buffer solutions.

0 3 . 1

Define Brønsted–Lowry base.

[1 mark]

A proton (H^+) acceptor.

0 3 . 2

Write an equation, including state symbols, for the reaction of sodium with water.

[1 mark]



0 3 . 3

0.150 g of sodium metal is carefully added to some water. After the reaction, more water is added to give a solution with a final volume of 2000 cm³

Calculate the pH of this solution at 25 °C

Give your answer to **two** decimal places.The ionic product of water, $K_w = 1.00 \times 10^{-14} \text{ mol}^2 \text{ dm}^{-6}$ at 25 °C

[4 marks]

$n(\text{Na}) = 0.150 \div 23.0 = 6.52 \times 10^{-3} \text{ mol} = n(\text{NaOH})$

$[\text{OH}^-] = 6.52 \times 10^{-3} \div 2.000 \text{ dm}^3 = 3.26 \times 10^{-3} \text{ mol dm}^{-3}$

$[\text{H}^+] = K_w \div [\text{OH}^-] = 1.00 \times 10^{-14} \div 3.26 \times 10^{-3}$

$[\text{H}^+] = 3.07 \times 10^{-12} \text{ mol dm}^{-3}$

$\text{pH} = -\log_{10}(3.07 \times 10^{-12}) = 11.51$

pH 11.51

0 3 . 4

The value of K_w is $1.47 \times 10^{-14} \text{ mol}^2 \text{ dm}^{-6}$ at 30°C

Calculate the pH of pure water at 30°C
Give your answer to **two** decimal places.

State why water is neutral, even though the pH at 30°C is **not** 7.00

[3 marks]

In pure water $[H^+] = [OH^-]$, so $[H^+] = \sqrt{K_w}$

$[H^+] = \sqrt{(1.47 \times 10^{-14})} = 1.21 \times 10^{-7} \text{ mol dm}^{-3}$

$\text{pH} = -\log_{10}(1.21 \times 10^{-7}) = 6.92$

pH 6.92

Reason why water is neutral *The concentrations of H^+ and OH^- ions are equal, $[H^+] = [OH^-]$.*

0 3 . 5

10.0 cm³ of 0.150 mol dm⁻³ sodium hydroxide solution are added to
10.0 cm³ of 0.205 mol dm⁻³ propanoic acid.

For propanoic acid, $\text{p}K_a = 4.87$

Calculate the pH of the buffer solution that is formed.
Give your answer to **two** decimal places.

[6 marks]

$n(\text{NaOH}) = 10.0 \times 0.150 \div 1000 = 1.50 \times 10^{-3} \text{ mol}$

$n(\text{propanoic acid}) = 10.0 \times 0.205 \div 1000 = 2.05 \times 10^{-3} \text{ mol}$

$n(\text{acid remaining}) = 2.05 \times 10^{-3} - 1.50 \times 10^{-3} = 5.50 \times 10^{-4} \text{ mol}$

$n(\text{sodium propanoate}) = 1.50 \times 10^{-3} \text{ mol}$

$K_a = 10^{-4.87} = 1.35 \times 10^{-5} \text{ mol dm}^{-3}$

$[H^+] = K_a \times [HA] \div [A^-] = 1.349 \times 10^{-5} \times (5.50 \times 10^{-4} \div 1.50 \times 10^{-3})$

$[H^+] = 4.95 \times 10^{-6} \text{ mol dm}^{-3}$

$\text{pH} = -\log_{10}(4.95 \times 10^{-6}) = 5.31$

pH 5.31

15

Turn over ►




0 4

This question is about Period 3 elements and their compounds.

0 4 . 1

Write an equation for the reaction between phosphorus and chlorine to make phosphorus(V) chloride.

[1 mark]



0 4 . 2

Write an equation for the reaction of phosphorus(V) chloride with water to make phosphoric(V) acid.


[1 mark]



0 4 . 3

Draw the displayed formula of phosphoric(V) acid.


[1 mark]



0 4 . 4

Write an equation to show how phosphoric(V) acid forms an acidic solution in water.

[1 mark]



Aluminium oxide is amphoteric.

0 4 . 5

Define amphoteric.

[1 mark]

It can act as both an acid and a base (it reacts with both acids and bases).

0 4 . 6

Write an equation to show how aluminium oxide can act as a base.

[1 mark]



0 4 . 7

Sulfur dioxide reacts with water to form an acid.

State the formula of the acid.

Suggest the approximate pH of a 0.1 mol dm^{-3} solution of this acid.

[2 marks]

Formula H_2SO_3

pH 2

8

Turn over for the next question

Turn over ►



0 5

This question is about ligands and complexes.

0 5 . 1

Transition metal complexes have different shapes.

Complete **Table 6** to name the shape of each complex.**[4 marks]****Table 6**


Complex	Name of shape
$[\text{CoCl}_4]^{2-}$	<i>tetrahedral</i>
$\text{Pt}(\text{NH}_3)_2\text{Cl}_2$	<i>square planar</i>
$[\text{Ag}(\text{NH}_3)_2]^+$	<i>linear</i>
$[\text{Cu}(\text{EDTA})]^{2-}$	<i>octahedral</i>

0 5 . 2

Ethanedioate ions ($\text{C}_2\text{O}_4^{2-}$) can act as bidentate ligands.

Draw the displayed formula of an ethanedioate ion.

Show on your diagram the lone pairs of electrons that form the co-ordinate bonds when the ion is acting as a ligand.

[2 marks]

0 5 . 3

When ethanedioate ions are added to an aqueous solution containing iron(II) ions, a ligand substitution reaction takes place.

Write an equation for the reaction that occurs when all six water ligands in the aqueous iron(II) complex are replaced by ethanedioate ions.

[1 mark]

0 5 . 4 1,2-diaminoethane is also a bidentate ligand.

When 1,2-diaminoethane is added to an aqueous solution containing copper(II) ions, incomplete substitution takes place to form the complex $[\text{Cu}(\text{NH}_2\text{CH}_2\text{CH}_2\text{NH}_2)_2(\text{H}_2\text{O})_2]^{2+}$

What type of isomerism does this complex show?

Tick (✓) **one** box.

[1 mark]

Cis-trans only

☐

Optical only

☐

Cis-trans and optical

☒

0 5 . 5 Haem is a complex that contains a multidentate ligand.

Identify the metal ion in this complex.

Explain how the complex allows oxygen to be transported in the body.

[2 marks]

Metal ion Fe^{2+}

Explanation Oxygen forms a co-ordinate bond to the Fe^{2+} ion (using a lone pair), and this bond is weak so the O_2 can be released where it is needed.

10

Turn over for the next question

Turn over ►



0 6

This question is about reactions of metal ions in aqueous solution.

Table 7 shows the observations from some test-tube reactions of an aqueous copper(II) salt.

Table 7

	Test	Observations
1	Add acidified BaCl ₂ (aq)	White precipitate
2	Add NaOH(aq)	Blue precipitate
3	Add excess HCl(aq)	Yellow-green solution

0 6

1

Use the observation from **Test 1** to identify the anion in the salt.

[1 mark]

sulfate ion, SO₄²⁻

0 6

2

Write an ionic equation for the reaction in **Test 2**.

[1 mark]

[Cu(H₂O)₆]²⁺ + 2OH⁻ → [Cu(H₂O)₄(OH)₂] + 2H₂O

0 6

3

The reaction in **Test 3** involves a change in the co-ordination number of the complex ion.

Write an equation for the reaction in **Test 3**.

Explain why there is a change in the co-ordination number.

[2 marks]

Equation

[Cu(H₂O)₆]²⁺ + 4Cl⁻ → [CuCl₄]²⁻ + 6H₂O

Explanation *The Cl⁻ ligand is larger than the H₂O ligand, so only four Cl⁻ ions fit around the Cu²⁺ ion (co-ordination number changes from 6 to 4).*



0 6 . 4 Table 8 shows two further test-tube reactions of the aqueous copper(II) salt.

Table 8

	Test	Observation
4	Add a few drops of $\text{NH}_3(\text{aq})$	<i>blue precipitate</i>
5	Add excess $\text{NH}_3(\text{aq})$	<i>precipitate dissolves to form a deep-blue solution</i>

Complete **Table 8** to state the observation in each test.

Give the formula of the copper-containing complex ion formed in **Test 5**.

[3 marks]

Formula of copper-containing complex ion $[\text{Cu}(\text{NH}_3)_4(\text{H}_2\text{O})_2]^{2+}$

An aqueous solution of sodium carbonate is added to separate samples of aqueous iron(II) sulfate and aqueous iron(III) chloride.

0 6 . 5 Describe what you would observe in the reaction with iron(II) sulfate.

Write an equation for the reaction with iron(III) chloride.

[2 marks]

Observation with iron(II) sulfate *green precipitate*

Equation with iron(III) chloride



0 6 . 6 Explain why aqueous iron(III) chloride is more acidic than aqueous iron(II) sulfate.

[2 marks]

Fe^{3+} has a higher charge and is smaller, so it has a higher charge density and is more polarising than Fe^{2+} . So it weakens the O-H bonds in the water ligands more and more H^+ ions are released (hydrolysis equilibrium lies further to the right).



0 7

Under some conditions, nitrate(V) ions (NO_3^-) can oxidise Mn^{2+} ions to MnO_2 and the nitrate(V) ions are reduced.

An experiment is used to find the oxidation state of nitrogen at the end of the reaction.

In this experiment, 24.0 cm^3 of $0.150 \text{ mol dm}^{-3}$ aqueous solution of Mn^{2+} ions react with exactly 15.0 cm^3 of $0.0960 \text{ mol dm}^{-3}$ aqueous solution of NO_3^- ions.

Write a half-equation for the oxidation of Mn^{2+} to MnO_2

Calculate the simplest whole-number reacting ratio of Mn^{2+} ions to NO_3^- ions.

Determine the oxidation state of N at the end of the reaction.

[5 marks]

Half-equation



Calculation

$$n(\text{Mn}^{2+}) = 24.0 \times 0.150 \div 1000 = 3.60 \times 10^{-3} \text{ mol}$$

$$n(\text{NO}_3^-) = 15.0 \times 0.0960 \div 1000 = 1.44 \times 10^{-3} \text{ mol}$$

$$\text{ratio} = 3.60 \times 10^{-3} \div 1.44 \times 10^{-3} = 2.5, \text{ so } \text{Mn}^{2+} : \text{NO}_3^- = 5 : 2$$

$$5\text{Mn}^{2+} \text{ lose } 5 \times 2 = 10 \text{ electrons, so } 2\text{NO}_3^- \text{ gain 10 electrons}$$

$$\text{each N gains 5 electrons: oxidation state } +5 \rightarrow 0$$

Reacting ratio

$\text{Mn}^{2+} : \text{NO}_3^-$

5 : 2

Oxidation state of N at the end of the reaction

0

5



Turn over for the next question

*Do not write
outside the
box*

**DO NOT WRITE ON THIS PAGE
ANSWER IN THE SPACES PROVIDED**


Turn over ►



0 8

Figure 3 shows the electrochemical cell used to measure the standard electrode potential for the Zn^{2+}/Zn electrode.

Figure 3



0 8

1

Identify a substance used in **W**.

Give the purpose of **W**.

[2 marks]

Substance potassium nitrate solution, $\text{KNO}_3(\text{aq})$

Purpose It allows ions to flow between the two half-cells, completing the circuit (without the solutions mixing).

0 8

2

Identify substances **X**, **Y** and **Z**.

[3 marks]

X platinum (platinum black electrode)

Y zinc

Z zinc nitrate solution, $1.00 \text{ mol dm}^{-3} \text{ Zn}^{2+}(\text{aq})$



Table 9 shows some electrode half-equations and their standard electrode potentials.

Table 9

Electrode half-equation	$E^\ominus / \text{V}$
$\frac{1}{2}\text{O}_2(\text{g}) + 2\text{H}^+(\text{aq}) + 2\text{e}^- \rightarrow \text{H}_2\text{O}(\text{l})$	+1.23
$\text{VO}_2^+(\text{aq}) + 2\text{H}^+(\text{aq}) + \text{e}^- \rightarrow \text{VO}^{2+}(\text{aq}) + \text{H}_2\text{O}(\text{l})$	+1.00
$\text{Ag}^+(\text{aq}) + \text{e}^- \rightarrow \text{Ag}(\text{s})$	+0.80
$\text{VO}^{2+}(\text{aq}) + 2\text{H}^+(\text{aq}) + \text{e}^- \rightarrow \text{V}^{3+}(\text{aq}) + \text{H}_2\text{O}(\text{l})$	+0.34
$\text{Pb}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Pb}(\text{s})$	-0.13
$\text{V}^{3+}(\text{aq}) + \text{e}^- \rightarrow \text{V}^{2+}(\text{aq})$	-0.26
$\text{Zn}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Zn}(\text{s})$	-0.76
$\text{V}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{V}(\text{s})$	-1.20

An excess of zinc reacts with an acidified solution containing aqueous VO_2^+ ions.

Several colour changes are seen as the reaction mixture is observed.

The final reaction mixture contains Zn^{2+} ions and V^{2+} ions.

0 8 . 3

Explain why there are several colour changes.

[2 marks]

Vanadium has several oxidation states, each with a different colour.

$E^\ominus$ of each vanadium half-equation is more positive than $E^\ominus(\text{Zn}^{2+}/\text{Zn}) = -0.76 \text{ V}$, so zinc reduces vanadium step by step from +5 to +4 to +3 to +2 ($\text{VO}_2^+ \rightarrow \text{VO}^{2+} \rightarrow \text{V}^{3+} \rightarrow \text{V}^{2+}$).

Question 8 continues on the next page

Turn over ►



Table 9 is repeated here.

Table 9

Electrode half-equation	$E^\ominus / \text{V}$
$\frac{1}{2}\text{O}_2(\text{g}) + 2\text{H}^+(\text{aq}) + 2\text{e}^- \rightarrow \text{H}_2\text{O}(\text{l})$	+1.23
$\text{VO}_2^+(\text{aq}) + 2\text{H}^+(\text{aq}) + \text{e}^- \rightarrow \text{VO}^{2+}(\text{aq}) + \text{H}_2\text{O}(\text{l})$	+1.00
$\text{Ag}^+(\text{aq}) + \text{e}^- \rightarrow \text{Ag}(\text{s})$	+0.80
$\text{VO}^{2+}(\text{aq}) + 2\text{H}^+(\text{aq}) + \text{e}^- \rightarrow \text{V}^{3+}(\text{aq}) + \text{H}_2\text{O}(\text{l})$	+0.34
$\text{Pb}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Pb}(\text{s})$	-0.13
$\text{V}^{3+}(\text{aq}) + \text{e}^- \rightarrow \text{V}^{2+}(\text{aq})$	-0.26
$\text{Zn}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Zn}(\text{s})$	-0.76
$\text{V}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{V}(\text{s})$	-1.20

0 8 . 4

The unreacted zinc is removed from the mixture.

This mixture is then left to stand in air.

A colour change is observed in the mixture that has been left to stand in air.

Use data from **Table 9** to explain this observation.

[2 marks]

V^{2+} ions are oxidised by oxygen in the air to V^{3+} (violet to green).

This happens because $E^\ominus(\text{O}_2/\text{H}_2\text{O}) = +1.23 \text{ V}$ is more positive than

$E^\ominus(\text{V}^{3+}/\text{V}^{2+}) = -0.26 \text{ V}$, so $\text{EMF} = +1.49 \text{ V}$.

$\frac{1}{2}\text{O}_2 + 2\text{H}^+ + 2\text{V}^{2+} \rightarrow 2\text{V}^{3+} + \text{H}_2\text{O}$

9

END OF QUESTIONS



There are no questions printed on this page

*Do not write
outside the
box*

**DO NOT WRITE ON THIS PAGE
ANSWER IN THE SPACES PROVIDED**



[illegible]

[illegible]

[illegible]

There are no questions printed on this page

*Do not write
outside the
box*

**DO NOT WRITE ON THIS PAGE
ANSWER IN THE SPACES PROVIDED**

Copyright information

For confidentiality purposes, all acknowledgements of third-party copyright material are published in a separate booklet. This booklet is published after each live examination series and is available for free download from www.oxfordaqaexams.org.uk.

Permission to reproduce all copyright material has been applied for. In some cases, efforts to contact copyright-holders may have been unsuccessful and Oxford International AQA Examinations will be happy to rectify any omissions of acknowledgements. If you have any queries please contact the Copyright Team.

Copyright © 2023 Oxford International AQA Examinations and its licensors. All rights reserved.

