

WJEC A-LEVEL CHEMISTRY • UNIT 2

CHEMISTRY



EXAM QUESTION PACK

2.3 THE WIDER IMPACT OF CHEMISTRY

SCORE

/141

PERCENTAGE

%

STEMUP

These questions belong to WJEC, not to us.

The examination questions, mark schemes and examiner comments in this pack were written and published by WJEC. They are past assessment materials, reproduced here so that students can revise from them.

What we do not own

STEMUP is not affiliated with WJEC and does not own, and makes no claim to own, any of the question, mark scheme or examiner report material in this pack. WJEC has not endorsed, approved or been involved in the production of it. WJEC and the qualification titles used here belong to their respective owners and appear for identification only.

Where to find the originals

WJEC publishes its past papers, mark schemes and examiner reports free of charge. Always check the current specification at wjec.co.uk before you revise, because content and wording can change between series.

If you hold the rights to any material in this pack and would like it amended or removed, contact admin@stemup.co.uk and we will act on it.

1. (a) Planners have to ensure a secure supply of energy in the future. It has been suggested that the use of fossil fuels should be reduced, the use of renewable energy increased and that energy efficiency should be greatly improved.

By considering both the benefits and the difficulties involved, discuss whether you think that these suggestions are realistic.

[4] QWC [1]

(i) I. State the general variations in this yield with temperature and pressure

[1]

II. Use the graphs to explain whether the reaction is endothermic or exothermic and whether there are more moles of gaseous products than reactants.

[4] QWC [1]

(ii) Normally the process is carried out at a temperature of around 900 °C.

Suggest why this temperature is used.

[2]


(iii) State the **type** of catalyst used.

[1]


(iv) Explain why there has been much research to find a better catalyst.

[2]

- (v) The next stage in the Ostwald process is to convert the nitric oxide to nitrogen dioxide.



Sketch on the axes below the energy profile for this reaction, clearly labelling the enthalpy change of reaction, ΔH . [2]




- (vi) Write an expression that connects the enthalpy change of a reaction, ΔH , with the activation energies of the forward (E_f) and reverse (E_b) reactions. [1]

Total [19]

2.

Hydrocarbons play an important role in our life today, both as fuels and as raw materials for the synthesis of a wide range of materials. Most hydrocarbons are isolated from crude oil, however there is increasing interest in alternative methods of obtaining these molecules.

- (a) One route to the production of hydrocarbons is the Fischer-Tropsch process, which uses hydrogen and carbon monoxide as starting materials to produce a range of molecules. The equation below shows the production of pentane, C_5H_{12} , by this route.



The enthalpies of formation of some of these substances are given in the table below.

Substance	Standard enthalpy of formation, $\Delta H_f^\ominus$ / kJ mol^{-1}
Hydrogen, $\text{H}_2(\text{g})$	0
Carbon monoxide, $\text{CO}(\text{g})$	-111
Water, $\text{H}_2\text{O}(\text{l})$	-286

- (i) State the temperature and pressure used as standard conditions. Give units for each. [2]

Temperature Pressure

- (ii) State why the standard enthalpy of formation for hydrogen gas is 0 kJ mol^{-1} . [1]

.....

.....

.....

- (iii) Use the values given to calculate the standard enthalpy of formation for pentane, $\text{C}_5\text{H}_{12}(\text{l})$, in kJ mol^{-1} . [3]

.....

.....

.....

.....

(b) The Fischer-Tropsch process uses a heterogeneous catalyst containing iron.

(i) State what is meant by the term *heterogeneous* in this context. [1]

.....

.....

(ii) Explain how a catalyst increases the rate of a chemical reaction. [2]

.....

.....

(iii) Chemical manufacturers consider catalysts to be a key part of production methods that have the minimum possible effect on the environment ('Green Chemistry'). Give **one** reason why the use of catalysts reduces the effect on the environment.[1]

.....

.....

.....

(iv) An alternative method of increasing the rate of a chemical reaction is to increase the temperature. Explain why temperature affects the rate of a chemical reaction. [3]
QWC [1]

.....

.....

.....

.....

- (c) One method of producing the hydrogen gas required for the Fischer-Tropsch process is to use the reversible reaction below.



- (i) State and explain the effect, if any, of increasing pressure on the yield of hydrogen gas produced at equilibrium. [2]

- (ii) State and explain the effect, if any, of increasing temperature on the yield of hydrogen gas produced at equilibrium. [2]

- (iii) This reaction uses a catalyst based on iron oxide. State the effect of using a catalyst on the position of equilibrium. [1]

Total [19]

3. (a) The combustion of fossil fuels containing sulfur impurities is known to cause acid rain. This acid rain can cause the erosion of marble statues as the calcium carbonate in them reacts with the acid in the rain.


Give **one** other problem caused by acid rain.

[1]

- (b) A chemist is developing coatings for marble that will slow down the rate of their erosion by acid rain. To compare different coatings he uses small model statues, all of which are the same size and shape as each other. He proposes to measure the rate of reaction by adding acid and measuring the volume of gas given off at set time intervals.

(i) Complete the diagram to show the apparatus that could be used to perform this experiment.

[1]




(ii) Explain why it is important that the model statues are the same size and shape as each other

[1]

(ii) State **two** other factors he will need to keep constant if he is to collect valid data.

[2]

(c) One gas that causes acid rain is sulfur dioxide. This gas is used to produce sulfur trioxide in the Contact Process. The reaction occurring is shown in the following equation.



(i) State and explain the effect of increasing pressure on the equilibrium yield of sulfur trioxide.

[2]

(ii) When the temperature is increased the rate at which equilibrium is reached is increased and the yield of sulfur trioxide is decreased.

I State whether this reaction is endothermic, exothermic or neither, giving a reason for your answer.

[2]

II Explain why increasing the temperature leads to an increase in the rate of reaction.

[3]

III To increase the rate of a reaction, a catalyst can be used. Give a **different** catalysed reaction and name the catalyst for this reaction.

[1]

- (d) Ethanoic acid, CH_3COOH , is one of the most familiar compounds used as a flavouring and preservative for food. Originally ethanoic acid was produced by oxidation of ethanol by bacteria in the presence of air (route A below). Today there are many other possible routes and three of these are shown as routes B, C and D below.

Route	Carbon-containing starting materials	Conditions	Overall equation	Atom economy
A	ethanol		$\text{C}_2\text{H}_5\text{OH} + \text{O}_2 \rightarrow \text{CH}_3\text{COOH} + \text{H}_2\text{O}$	76.9%
B	methanol, carbon monoxide	150 °C, 30 atm	$\text{CH}_3\text{OH} + \text{CO} \rightleftharpoons \text{CH}_3\text{COOH}$	100.0%
C	butane	150 °C, 55 atm	$2\text{C}_4\text{H}_{10} + 5\text{O}_2 \rightarrow 4\text{CH}_3\text{COOH} + 2\text{H}_2\text{O}$	87.0%
D	sugars		$\text{C}_6\text{H}_{12}\text{O}_6 \rightarrow 3\text{CH}_3\text{COOH}$	

- (i) State the atom economy of route D for production of ethanoic acid.

[1]

- (ii) Route B is the route most commonly used for producing ethanoic acid today for both financial and *Green Chemistry* reasons. Apply the principles of *Green Chemistry* to the information above to give **two** reasons why route B is favoured over route C.

[2]

- _____
- _____

- (iii) Route B uses a homogeneous catalyst. State what effect the catalyst will have on the position of this equilibrium.

[1]

(Total 17)

4. (a) The vast majority of motor vehicles worldwide are powered by petrol or diesel which come from crude oil. Give **two** reasons why we cannot rely indefinitely on oil as a source of transport fuel.

[2]

(b) Many vehicle manufacturers around the world have made the development of alternative fuels a priority. One such fuel being studied is hydrogen.

Its main advantage is that the only waste product is water, however hydrogen does not occur naturally on Earth. It is produced by passing an electric current through water.

(i) A leading car manufacturer said:

“Cars powered by hydrogen will be pollution-free”

Give **two** reasons why this is not necessarily true.

[2] QWC [1]

(ii) A spokesperson for a safety group said,

“Hydrogen can burn explosively. It must not be used in cars unless it is 100 % safe”.

State, giving a reason, whether you agree with this.

[1]

(c) The first line in the visible atomic emission spectrum for hydrogen has a wavelength of 656 nm, while that for helium has a wavelength of 707 nm.

State, giving a reason, which line has:

(i) the higher frequency,

[1]

(ii) the higher energy.

[1]

(d) The first ionisation energy of helium is 2370 kJ mol^{-1} while that of neon is 2080 kJ mol^{-1} . Explain why neon has a lower first ionisation energy than helium.

[2]

(e) Another noble gas is radon. Its more stable isotope ^{222}Rn has a half-life of 3.8 days, decays by α -emission and is responsible for the majority of the public exposure to ionising radiation.

(i) Give the symbol and mass number of the atom formed by the loss of one α -particle from an atom of ^{222}Rn .

[1]


(ii) Explain why doctors are concerned that an over-exposure to radon may cause lung cancer.

[1]

(Total 12)

5.

- (a) Nitrogen(I) oxide is a colourless gas that reacts with hydrogen to give nitrogen and water.



- (i) State why the standard enthalpy of formation of both hydrogen and nitrogen gases is 0 kJ mol^{-1} . [1]


- -----
- (ii) Calculate the standard enthalpy of formation of nitrogen(I) oxide in kJ mol^{-1} .
(You should assume that the standard enthalpy of formation of water is -286 kJ mol^{-1}) [2]

Standard enthalpy of formation = ... kJ mol^{-1}

- (b) A new method for producing phenol, $\text{C}_6\text{H}_5\text{OH}$, is by reacting benzene, C_6H_6 , with nitrogen(I) oxide at 400°C in the presence of a suitable catalyst.



- (i) Sketch the energy profiles for the catalysed and uncatalysed reactions using the axes shown below.
Label your profiles as *catalysed* and *uncatalysed*. [2]



- (ii) A pilot-scale plant used 156 kg of benzene ($M_r = 78$) to produce phenol ($M_r = 94$).

I Calculate the number of moles of benzene used. [1]

Moles of benzene = ... mol


- II The yield of phenol was 95 %. Using your answer to I and the equation below (or another suitable method), calculate the mass of phenol obtained. Show your working. [3]



Mass of phenol = ... kg

- (iii) Study the short account below, which gives more detail about this process.

The process to make phenol is carried out in the gas phase and uses a solid zeolite catalyst. The operating temperature is around 400 °C.



The reactants are the hydrocarbon benzene and nitrogen(I) oxide, which is a potent greenhouse gas. The nitrogen(I) oxide is obtained from another process, where it is produced as an undesirable side product.

Use the account and the equation to comment on the environmental and *Green Chemistry* advantages of this process. A reference to the yield is not required. [4]

QWC [1]

Total [14]

6.

- (a) Potassium hydroxide contains potassium ions, K^+ .
Give the electron configuration of a potassium **atom** and use this to explain why most potassium compounds contain the potassium ion. [2]

- (b) Michael was asked to make 250 cm^3 of a solution of potassium hydroxide and to record the maximum rise in temperature that occurred as it dissolved.
He measured 250 cm^3 of water in a glass beaker and then added 7.01 g (0.125 mol) of solid potassium hydroxide to this, with stirring.
He noticed that the temperature rose from 20.2°C to a maximum of 25.0°C .

- (i) Calculate the molar enthalpy change of solution of potassium hydroxide by use of the formula

$$\Delta H = - \frac{mc\Delta T}{n}$$

where m = mass of the solvent in grams (assume 1 cm^3 has a mass of 1 g)
 c = $4.2\text{ J g}^{-1}^\circ\text{C}^{-1}$
 ΔT = change in temperature of the solution
 n = number of moles of the solute
 ΔH = molar enthalpy change of solution

You should show the **units** in your answer. [3]

$$\Delta H = \dots\dots\dots$$

(ii) Michael's measurements produced a value for the enthalpy of solution of potassium hydroxide that was different to the literature value.

Use the information given to suggest and explain **two** factors that might produce a different result.

[2]

1.
.....
2.
.....

(c) Solid potassium hydroxide can be used in analysis to find the percentage of carbon dioxide present in a mixture of gases. The equation for the reaction that occurs is given below.



2.0 m³ of a gas mixture was passed through potassium hydroxide. Analysis showed that 1.50 mol of potassium carbonate had been formed.

(i) State the number of moles of carbon dioxide necessary to produce 0.050 mol of potassium carbonate. [1]

(ii) Calculate the volume of carbon dioxide that produced 0.050 mol of potassium carbonate. [1]

[1 mol of carbon dioxide has a volume of 24.0 dm³ under these conditions]

Volume of carbon dioxide = ... dm³

(iii) Calculate the percentage of carbon dioxide in the gas mixture, in terms of volume. [2]

[1 dm³ = 0.001 m³]

(d) Scientists have commented that 'an increase in the amount of carbon dioxide dissolved in seawater will cause problems for animals whose shells are composed of calcium carbonate'.



Use the equation above to help you discuss the problem that is caused for these animals by this increase in carbon dioxide concentration.

[3] QWC [1]

(Total 15)

7.

(a) State what is meant by the term *standard molar enthalpy change of formation*. [2]

(b) (i) Write an equation to represent the standard molar enthalpy change of formation, $\Delta H_f^\ominus$, of $\text{H}_2\text{O}(\text{g})$. [1]

(ii) The standard molar enthalpy change of formation, $\Delta H_f^\ominus$, of $\text{H}_2\text{O}(\text{g})$ is -242 kJ mol^{-1} . Using this value and the average bond enthalpies given in the table below, calculate the average bond enthalpy of the O—H bond in H_2O . [2]

Bond	Average bond enthalpy / kJ mol^{-1}
H—H	436
O=O	496

- (c) Hydrogen has been proposed as a possible alternative to petrol as a fuel for cars. One suggestion is to store the hydrogen in the car as solid magnesium hydride, MgH_2 , and generate it as required by heating.

(i) I Give **one** advantage of using hydrogen in place of petrol as a fuel for cars. [1]

.....
.....

II Give **one** advantage of storing the fuel in the car in the form of magnesium hydride rather than hydrogen gas. [1]

.....
.....

- (ii) One possible disadvantage of using magnesium hydride arises from its reaction with water.



Suggest why magnesium hydride's reaction with water could be a problem. [1]

.....
.....


- (iii) The fuel tank of one type of hydrogen-powered car holds 70 kg of magnesium hydride.

Calculate the volume of hydrogen gas, measured at room temperature and pressure, which would be produced if this amount of magnesium hydride reacted with water. [3]

[1 mol of gas molecules occupies 24 dm^3 at room temperature and pressure]

Volume of hydrogen gas = ... dm^3

- (d) Methanol can be produced industrially by passing carbon monoxide and hydrogen over a catalyst at high temperatures and pressures.



(i) State how the equilibrium yield of methanol is affected by an increase in temperature and in pressure.

[1]

(ii) Explain your answer to part (i).

[2]

(e) Many catalysts are very expensive but their use does allow the chemical industry to operate more profitably. Explain why the use of catalysts provides economic and environmental benefits.

[3] QWC [1]

(Total 18)

2.

- (a) (i) Temperature: 298K / 25°C (1) Pressure: 1 atm / 101.325 kPa or 100 kPa (1) [2]
- (ii) Hydrogen gas is an element in its standard state [1]
- (iii) $\Delta H = \Delta H_f(C_5H_{12}) + 5 \Delta H_f(H_2O) - 5 \Delta H_f(CO) - 11\Delta H_f(H_2)$ (1)
- $\Delta H_f(C_5H_{12}) = -1049 - 5(-286) + 5(-111)$ (1)
- $\Delta H_f(C_5H_{12}) = -174 \text{ kJ mol}^{-1}$ (1) [3]
- (b) (i) Catalyst in different state to reactants [1]
- (ii) Catalysts provide an alternative route (1) with a lower activation energy (1) [2]
- (iii) Lower temperature or less time so less energy needed / Can make alternative production method possible with sustainable starting materials or less waste products [1]
- (iv) At higher temperatures particles have more energy (1)
- More collisions have energy above activation energy (1)
- (Can obtain these two marks from correctly labelled Boltzmann energy distribution plot with two temperature lines (1) and Activation energy (1))*
- Successful collisions occur more frequently (1) – 3 max [3]
- QWC: selection of a form and style of writing appropriate to purpose and to complexity of subject matter* [1]
- (c) (i) No effect (1)
- Same number of (gas) molecules on both sides of reaction (1) [2]
- (ii) Lower yield of hydrogen (1)
- Reaction shifts in endothermic direction to (try to counteract increase in temperature) (1) [2]
- (iii) No effect [1]

Total [19]

3.

(a)	killing marine life / killing trees	[1]
(b)	(i) either gas syringe or inverted burette attached to sealed vessel	[1]
	(ii) different surface area would affect rate of reaction	[1]
	(iii) concentration / volume / nature of acid (1)	
	temperature (1)	[2]
(c)	(i) increasing pressure will shift the reaction to side with fewer gas molecules (1)	
	increasing yield of SO_3 (1) – reason must be given	[2]
	(ii) I increasing temperature shifts equilibrium in endothermic direction (1)	
	as SO_3 yield is decreased forward reaction must be exothermic (1)	[2]
	II increasing temperature increases energy of particles (1)	
	more collisions have energy above activation energy (1)	
	successful collisions occur more frequently (1)	
	<i>can gain first two points from labelled Boltzmann distribution curve</i>	[3]
	III e.g. iron in production of ammonia or any valid example	[1]
(d)	(i) atom economy = 100%	[1]
	(ii) any two points from:	
	lower pressure used in B (1)	
	methanol is a renewable starting material (1)	
	higher atom economy in B or less waste in B (1)	
	[ignore reference to cost]	
	2 max	[2]
	(iii) no effect on position of equilibrium	[1]
Total		[17]

4.

- (a) Oil is non-renewable / will run out (1)
Contribution of CO₂ to global warming (1)
Oil has other important uses (1) [2]
(Maximum 2 marks)
- (b) (i) Power stations / fossil fuels used to generate the electricity needed to make H₂ (1)
Resulting in CO₂ formation (global warming) / acid rain (1)
Manufacture of car produces pollution (1) [2]
(Maximum 2 marks)
QWC Legibility of text; accuracy of spelling, punctuation and grammar, clarity of meaning [1]
- (ii) Disagree, no fuel is 100% safe / petrol can burn explosively
(Accept agree if valid reason given e.g. in terms of lives being lost) [1]
- (c) (i) Hydrogen since frequency is inversely proportional to wavelength / smaller wavelength [1]
- (ii) Hydrogen since energy is proportional to frequency / greater frequency / $E = hf$ [1]
- (d) In Ne greater shielding of *outer* electron (1) outweighs larger nuclear charge (1) / He has greater effective nuclear charge (1) / He *outer* electron closer to nucleus (1)
- max 1 if no reference to *outer* electron [2]
(Maximum 2 marks)
- (e) (i) ²¹⁸Po [1]
- (ii) Since radon is a gas / inhaled, α particles will be given off in the lungs (which may cause cancer) [1]

Total [12]


5.

(a) (i) They are both elements in their standard states. [1]

(ii) $\Delta H = \sum \Delta H_f \text{ products} - \sum \Delta H_f \text{ reactants}$ (1)
 $= (-286 + 0) - (-368 + 0)$
 $= -286 + 368 = (+)82 \text{ (kJ mol}^{-1}\text{)}$ (1) [2]

or by a cycle where correct cycle drawn (1) correct answer (1)

(b) (i)



exothermic profile drawn (1)
 uncatalysed / catalysed line labelled (1) [2]

(ii) I number of moles of benzene = 2000 [1]

II mole ratio is 1 : 1 (1)

$$\therefore \text{moles of phenol produced} = \frac{2000 \times 95}{100} = 1900 \text{ (1)}$$

$$\text{mass} = M_r \times \text{number of moles} = 94 \times 1900 = 178.6 / 179 \text{ kg (1)}$$

alternatively

78 (g / kg) of benzene gives 94 (g / kg) of phenol (1)

$\therefore$ 1 (g / kg) of benzene gives $94/78$ (g / kg) of phenol

$\therefore$ 156 (kg) of benzene gives $94 \times 156/78$ (kg) of phenol = 188 (kg) (1)

but 95% yield $\therefore \frac{188 \times 95}{100} = 178.6 / 179 \text{ (kg)} \text{ (1)} \quad [3]$

(iii) Look for at least four relevant positive points [4] e.g.

- the process uses a (heterogeneous) catalyst, which can easily be separated from the gaseous products (thus saving energy)
- the only other product of the reaction is gaseous nitrogen, which is non-toxic / safe / not a harmful product
- the process uses nitrogen(I) oxide which is used up, rather than being released into the atmosphere from the other process (and causing global warming)
- the process is exothermic and the heat produced can be used elsewhere
- a relatively moderate operating temperature reduces overall costs
- high atom economy

Legibility of text; accuracy of spelling, punctuation and grammar; clarity of meaning QWC [1]

Total [14]

6. (a) $K \rightarrow 1s^2 2s^2 2p^6 3s^2 3p^6 4s^1$ (1)
There is one outer electron and the loss of this electron gives a stable potassium ion with a full outer shell/ ion more stable than the atom (1) [2]
- (b) (i) $\Delta T = 4.8^\circ\text{C}$ (1)
$$\Delta H = \frac{-250 \times 4.2 \times 4.8}{0.125} = -40320 \text{ J mol}^{-1} / -40.3 \text{ kJ mol}^{-1}$$
 (2) [3]
✓ for negative sign
✓ correct value with relevant units
- (ii) e.g. The volume used was not precise in measurement as the readings on a beaker are only approximate (1)
The experiment was performed in a beaker and this was not insulated and heat was lost to the surroundings (1) [2]

there may be other acceptable answers here, for example based on slow dissolving
- (c) (i) 0.050 [1]
(ii) $(0.050 \times 24.0) = 1.20 \text{ (dm}^3\text{)}$ [1]
(iii) $\% \text{ v/v} = \frac{1.20 \times 0.001 \times 100}{2}$ (1) = 0.06 (1) [2]
- (d) An increase in the concentration of (aqueous) carbon dioxide causes the position of equilibrium to move to the right. (1)
This causes calcium carbonate to become aqueous calcium (and hydrogencarbonate) ions / dissolve (1)
weakening shells / causing difficulty in formation of shells (1) [3]

Organisation of information clearly and coherently; using specialist vocabulary where appropriate QWC [1]

Total [15]

7.

- (a) Enthalpy change when one mole of a compound is formed from its (constituent) elements (1)
in their standard states / under standard conditions (1) [2]
- (b) (i) $\text{H}_2 + \frac{1}{2}\text{O}_2 \rightarrow \text{H}_2\text{O}$ [1]
- (ii) $-242 = 436 + 248 - 2(\text{O}-\text{H})$ (1)
 $2(\text{O}-\text{H}) = 926$
 $\text{O}-\text{H} = 463 \text{ kJ mol}^{-1}$ (1) [2]
- (c) (i) I. Burning hydrogen will not produce CO_2 (or SO_2) as pollutants [1]
II. Hydrogen is very flammable, storing as MgH_2 is safer / MgH_2 is solid therefore volume occupied by given amount of hydrogen is less [1]
- (ii) If the MgH_2 is not kept dry, hydrogen will be formed and there could be a potential explosion [1]
- (iii) Moles $\text{MgH}_2 = \frac{70000}{26.32} = 2659.6$ (2660) (1)
Moles $\text{H}_2 = 5319.2$ (5320) (1)
Volume $\text{H}_2 = 1.28 \times 10^5 \text{ dm}^3$ (1) [3]
- (d) (i) An increase in temperature would decrease the yield and an increase in pressure would increase the yield [1]
- (ii) Forward reaction is exothermic so equilibrium shifts to the left as temperature is increased (1)
More gaseous moles on the l.h.s. so equilibrium shifts to the right as pressure is increased (1) [2]
- (e) Lower temperatures can be used (1)
Energy costs saved (1)
More product can be made in a given time (so more can be sold) (1)
Enable reactions to take place that would be impossible otherwise (1)
Less fossil fuels burned to provide energy (so less CO_2 formed) (1)
(any 3 of above) [3]
- QWC Legibility of text; accuracy of spelling, punctuation and grammar, clarity of meaning* [1]

Total [18]