GCE A LEVEL

1305U60-1

S23-1305U60-1

WEDNESDAY, 14 JUNE 2023 – AFTERNOON

FURTHER MATHEMATICS – A2 unit 6
FURTHER MECHANICS B

1 hour 45 minutes

1305U601
01

ADDITIONAL MATERIALS

In addition to this examination paper, you will need:

- a WJEC pink 16-page answer booklet;
- a Formula Booklet;
- a calculator.

INSTRUCTIONS TO CANDIDATES

Use black ink or black ball-point pen.

Answer **all** questions.

Take g as $9{\cdot}8\,\text{ms}^{-2}$.

Sufficient working must be shown to demonstrate the **mathematical** method employed.

Answers without working may not gain full credit.

Unless the degree of accuracy is stated in the question, answers should be rounded appropriately.

INFORMATION FOR CANDIDATES

The maximum mark for this paper is 80.

The number of marks is given in brackets at the end of each question or part-question.

You are reminded of the necessity for good English and orderly presentation in your answers.

© WJEC CBAC Ltd.

stemup.co.uk

Additional Formulae for 2023

Laws of Logarithms

$$\log_a x + \log_a y \equiv \log_a (xy)$$

$$\log_a x - \log_a y \equiv \log_a \left(\frac{x}{y}\right)$$

$$k \log_a x \equiv \log_a \left(x^k\right)$$

Sequences

General term of an arithmetic progression:

$$u_n = a + (n-1)d$$

General term of a geometric progression:

$$u_n = ar^{n-1}$$

Mensuration

For a circle of radius, r, where an angle at the centre of θ radians subtends an arc of length s and encloses an associated sector of area A :

$$s = r\theta \qquad A = \frac{1}{2}r^2\theta$$

Calculus and Differential Equations

Differentiation

Function	Derivative
$f(x)g(x)$	$f'(x)g(x) + f(x)g'(x)$
$f\left(g(x)\right)$	$f'\left(g(x)\right)g'(x)$

Integration

Function	Integral
$f'\left(g(x)\right)g'(x)$	$f\left(g(x)\right) + c$


Area under a curve $= \int_a^b y \, \mathrm{d}x$

© WJEC CBAC Ltd.

stemup.co.uk

Reminder: Sufficient working must be shown to demonstrate the **mathematical** method employed.

1. The diagram shows a uniform rod AB, of length 8 m and mass 23 kg, in limiting equilibrium with its end A on rough horizontal ground and point C resting against a smooth fixed cylinder. The rod is inclined at an angle of 30° to the ground.



The coefficient of friction between the ground and the rod is $\frac{2}{3}$.


(a) Calculate the magnitude of the normal reaction at C and the magnitude of the normal reaction to the ground at A. [8]

(b) Find the length AC. [4]

(c) Suppose instead that the rod is **non-uniform** with its centre of mass closer to A than to B. **Without carrying out any further calculations**, state whether or not this will affect your answers in part (a). Give a reason for your answer. [1]

2. *You are given that the centre of mass of a uniform solid cone of height h and base radius r is at a height of $\frac{1}{4}h$ above its base.*

The diagram shows a solid conical frustum. It is formed by taking a uniform right circular cone, of base radius $3x$ and height $6y$, and removing a smaller cone, of base radius x, with the same vertex.




Show that the distance of the centre of mass of the frustum from its base along the axis of symmetry is $\frac{18}{13}y$. [7]

TURN OVER

© WJEC CBAC Ltd.

stemup.co.uk

3. The vertical motion of a point on the surface of the water in a certain harbour may be modelled as Simple Harmonic Motion about a mean level. The diagram shows that, on a particular day, the depth of water in the harbour at low tide is 2 m and the depth of the water in the harbour at high tide is 10 m. The table below shows the times of high and low tides on this day.



Tidal Times		
High/Low	Time	Depth (metres)
Low Tide	5 a.m.	2
High Tide	11 a.m.	10
Low Tide	5 p.m.	2
High Tide	11 p.m.	10

(a) Write down the period and amplitude of the motion. [2]

(b) Let x m denote the height of water above mean level t hours after 5 a.m. Find an expression for x in terms of t. [3]

(c) The depth of water must be at least 4 m for boats to safely use the harbour. Determine the earliest time, after low tide at 5 a.m., at which boats can safely leave the harbour and hence find the latest possible time of return before the next low tide. [4]


(d) Calculate the rate at which the level of water is falling at 2 p.m. [4]

© WJEC CBAC Ltd.

stemup.co.uk

4. The diagram shows three **light** rods *AB*, *BC* and *CA* rigidly joined together so that *ABC* is a right-angled triangle with $AB = 45\,\text{cm}$, $AC = 28\,\text{cm}$ and $C\hat{A}B = 90°$. The rods support a uniform lamina, of density $2m\,\text{kg/cm}^2$, in the shape of a quarter circle *ADE* with radius 12 cm and centre at the vertex *A*. Three particles are attached to *BC*: one at *B*, one at *C* and one at *F*, the midpoint of *BC*.

The masses at *C*, *F* and *B* are $50m\,\text{kg}$, $30m\,\text{kg}$ and $20m\,\text{kg}$ respectively.



(a) Calculate the distance of the centre of mass of the system from

(i) *AC*,

(ii) *AB*. [11]

(b) When the system is freely suspended from a point *P* on *AC*, it hangs in equilibrium with *AB* vertical. Write down the length *AP*. [1]

(c) When the system is freely suspended from a point *Q* on *AD*, it hangs in equilibrium with *QB* making an angle of 60° with the vertical. Find the distance *AQ*. [3]

TURN OVER

© WJEC CBAC Ltd.

stemup.co.uk

5. In this question, $\mathbf{i}$ and $\mathbf{j}$ represent unit vectors due east and due north respectively.

Two smooth spheres *P* and *Q*, of equal radii, are moving on a smooth horizontal surface. The mass of *P* is 2 kg and the mass of *Q* is 6 kg. The velocity of *P* is $(8\mathbf{i} - 6\mathbf{j})\,\text{ms}^{-1}$ and the velocity of *Q* is $(4\mathbf{i} + 10\mathbf{j})\,\text{ms}^{-1}$. At a particular instant, *Q* is positioned 12 m east and 48 m south of *P*.

(a) Prove that *P* and *Q* will collide. [4]

At the instant the spheres collide, the line joining their centres is parallel to the vector $\mathbf{j}$. Immediately after the collision, sphere *Q* has speed $5\,\text{ms}^{-1}$.

(b) Determine the coefficient of restitution between the spheres and hence calculate the velocity of *P* immediately after the collision. [9]

(c) Find the magnitude of the impulse required to stop sphere *P* after the collision. [3]

© WJEC CBAC Ltd.

stemup.co.uk

6. The diagram on the left shows a train of mass 50 tonnes approaching a buffer at the end of a straight horizontal railway track. The buffer is designed to prevent the train from running off the end of the track. The buffer may be modelled as a light horizontal spring *AB*, as shown in the diagram on the right, which is fixed at the end *A*. The train strikes the buffer so that *P* makes contact with *B* at $t = 0$ seconds.

While *P* is in contact with *B*, an additional resistive force of $250\,000v$ N will oppose the motion of the train, where v ms^{-1} is the speed of the train at time t seconds. The spring has natural length 1 m and modulus of elasticity 312 500 N. At time t seconds, the compression of the spring is x metres.



(a) Show that, while *P* is in contact with *B*, x satisfies the differential equation

$$4\frac{\mathrm{d}^2x}{\mathrm{d}t^2} + 20\frac{\mathrm{d}x}{\mathrm{d}t} + 25x = 0\,. \qquad [4]$$

(b) Given that, when *P* first makes contact with *B*, the speed of the train is U ms^{-1}, find an expression for x in terms of U and t. [8]

(c) When the train comes to rest, the compression of the buffer is 0·3 m. Determine the speed of the train when it strikes the buffer. [3]

(d) State which type of damping is described by the motion of *P*. Give a reason for your answer. [1]

END OF PAPER


© WJEC CBAC Ltd.

stemup.co.uk

Surname	StemUp Model Solution
Answer booklet	Write the question number in the left-hand margin at the start of each answer.

Question number

1 (a)



Resolving vertically: $S\cos30° + R = 23g$

Resolving horizontally: $F = S\sin30°$

Limiting friction: $F = \frac{2}{3}R \Rightarrow \frac{2}{3}R = \frac{1}{2}S \Rightarrow R = \frac{3}{4}S$

$\frac{\sqrt{3}}{2}S + \frac{3}{4}S = 23g = 225.4$

$S = 225.4 \div \left(\frac{\sqrt{3}}{2} + \frac{3}{4}\right) = 139.478...$

$R = \frac{3}{4} \times 139.478... = 104.608...$

Normal reaction at C = 139.5 N, normal reaction at A = 104.6 N

1 (b)

Moments about A: $S \times AC = 23g \times 4\cos30°$

$139.478... \times AC = 225.4 \times 4\cos30° = 780.808...$

$AC = \frac{780.808...}{139.478...} = 5.598... \therefore AC = 5.60\ m$

1 (c)

No. Both reactions remain the same: in (a) only resolving was used (no moments), so the answers do not depend on where the centre of mass is.

2

Small cone has radius x so, by similar cones, its height is 2y.

Its base is 6y − 2y = 4y above the base of the frustum.

Question number

Shape	Volume (mass)	c.o.m. above base
Large cone	$\frac{1}{3}\pi(3x)^2(6y) = 18\pi x^2 y$	$\frac{1}{4}(6y) = \frac{3y}{2}$
Small cone	$\frac{1}{3}\pi x^2(2y) = \frac{2}{3}\pi x^2 y$	$4y + \frac{1}{4}(2y) = \frac{9y}{2}$
Frustum	$18\pi x^2 y - \frac{2}{3}\pi x^2 y = \frac{52}{3}\pi x^2 y$	$\bar{y}$

Moments about the base:

$\frac{52}{3}\pi x^2 y \times \bar{y} = 18\pi x^2 y \times \frac{3y}{2} - \frac{2}{3}\pi x^2 y \times \frac{9y}{2}$

$\frac{52}{3}\bar{y} = 27y - 3y = 24y$

$\bar{y} = 24y \times \frac{3}{52} = \frac{72}{52}y = \frac{18}{13}y$ as required

3 (a) Period = 2 × (11 a.m. − 5 a.m.) = 12 hours

Amplitude = $\frac{10-2}{2} = 4$ m

3 (b) $\omega = \frac{2\pi}{12} = \frac{\pi}{6}$

At $t = 0$ it is low tide, so $x = -4$ when $t = 0$

$x = -4\cos\left(\frac{\pi}{6}t\right)$

3 (c) Mean level is 6 m, so depth 4 m ⇒ $x = -2$

$-2 = -4\cos\left(\frac{\pi}{6}t\right) \Rightarrow \cos\left(\frac{\pi}{6}t\right) = \frac{1}{2}$

$\frac{\pi}{6}t = \frac{\pi}{3}$ or $\frac{5\pi}{3} \Rightarrow t = 2$ or $t = 10$

Earliest time to leave: 7 a.m. Latest time to return: 3 p.m.

3 (d) $v = \frac{dx}{dt} = 4 \times \frac{\pi}{6}\sin\left(\frac{\pi}{6}t\right) = \frac{2\pi}{3}\sin\left(\frac{\pi}{6}t\right)$

At 2 p.m., $t = 9$: $v = \frac{2\pi}{3}\sin\left(\frac{9\pi}{6}\right) = \frac{2\pi}{3} \times (-1) = -\frac{2\pi}{3}$

Water level is falling at $\frac{2\pi}{3}$ m/hour (= 2.09 m/hour)

4 (a) Lamina mass = $\frac{1}{4}\pi(12)^2 \times 2m = 72\pi m$

Lamina c.o.m. is $\frac{4r}{3\pi} = \frac{4(12)}{3\pi} = \frac{16}{\pi}$ cm from both AC and AB

Question number

	Mass	Distance from AC	Distance from AB
Lamina	$72\pi m$	$\frac{16}{\pi}$	$\frac{16}{\pi}$
C	$50m$	0	28
F	$30m$	22.5	14
B	$20m$	45	0
System	$(72\pi + 100)m$	$\bar{x}$	$\bar{y}$

(i) Moments about AC:

$72\pi(\frac{16}{\pi}) + 50(0) + 30(22.5) + 20(45) = (72\pi + 100)\bar{x}$

$1152 + 675 + 900 = 2727 = (72\pi + 100)\bar{x}$

$\bar{x} = \frac{2727}{72\pi + 100} = 8.3600... = 8.36$ cm from AC

(ii) Moments about AB:

$72\pi(\frac{16}{\pi}) + 50(28) + 30(14) + 20(0) = (72\pi + 100)\bar{y}$

$1152 + 1400 + 420 = 2972 = (72\pi + 100)\bar{y}$

$\bar{y} = \frac{2972}{72\pi + 100} = 9.1111... = 9.11$ cm from AB

4 (b) G is vertically below P with AB vertical, so $AP = \bar{y} = 9.11$ cm

4 (c) Hanging in equilibrium, G is vertically below Q, so angle GQB = 60°

$\tan 60° = \frac{\bar{y}}{\bar{x} - AQ} \Rightarrow \bar{x} - AQ = \frac{9.1111...}{\sqrt{3}} = 5.2603...$

$AQ = 8.3600... - 5.2603... = 3.0997... \therefore AQ = 3.10$ cm

5 (a) $r_P = (8i - 6j)t$ $r_Q = (12i - 48j) + (4i + 10j)t$

Collide if $r_P = r_Q$ for the same value of t:


i: $8t = 12 + 4t \Rightarrow t = 3$

j: $-6t = -48 + 10t \Rightarrow 16t = 48 \Rightarrow t = 3$

Both components give $t = 3$, so P and Q collide (when $t = 3$ s).

Question number

5 (b)



Line of centres is parallel to j, so the i components are unchanged.

After: Q has velocity $4i + v_Q j$, P has velocity $8i + v_P j$

Speed of Q: $\sqrt{(4^2 + v_Q^2)} = 5 \Rightarrow v_Q = \pm 3$

Conservation of momentum (j): $2(-6) + 6(10) = 2v_P + 6v_Q$

$48 = 2v_P + 6v_Q$: $v_Q = 3 \Rightarrow v_P = 15$; $v_Q = -3 \Rightarrow v_P = 33$

Restitution (j): $v_Q - v_P = -e(10 - (-6)) = -16e$

$v_Q = 3$: $3 - 15 = -16e \Rightarrow e = \frac{3}{4}$

$v_Q = -3$: $-3 - 33 = -16e \Rightarrow e = \frac{9}{4} > 1$, impossible, reject

$\therefore e = \frac{3}{4}$ and velocity of P after collision = $(8i + 15j)$ m s^{-1}

5 (c)

I = change in momentum of P = $2(0 - (8i + 15j)) = -16i - 30j$

$|I| = \sqrt{(16^2 + 30^2)} = \sqrt{1156} = 34$ Ns

6 (a)

Tension (thrust) in spring: $T = \frac{\lambda x}{l} = \frac{312500x}{l} = 312500x$

N2L in the direction of increasing x (direction of motion):

$-312500x - 250000v = 50000a$, $v = \frac{dx}{dt}$, $a = \frac{d^2x}{dt^2}$

$50000\frac{d^2x}{dt^2} + 250000\frac{dx}{dt} + 312500x = 0 \quad (\div 12500)$

$4\frac{d^2x}{dt^2} + 20\frac{dx}{dt} + 25x = 0$ as required

6 (b)

Auxiliary equation: $4m^2 + 20m + 25 = 0 \Rightarrow (2m + 5)^2 = 0$

$m = -\frac{5}{2}$ (repeated root)

General solution: $x = (At + B)e^{-5t/2}$

$t = 0, x = 0 \Rightarrow B = 0$

Question number

$\frac{dx}{dt} = Ae^{-5t/2} - \frac{5}{2}(At + B)e^{-5t/2}$

$t = 0, \frac{dx}{dt} = U \Rightarrow U = A - \frac{5}{2}B \Rightarrow A = U$

$\therefore x = Ute^{-5t/2}$

6 (c) $v = \frac{dx}{dt} = Ue^{-5t/2}(1 - \frac{5}{2}t) = 0$ at rest $\Rightarrow t = \frac{2}{5}$

$x = 0.3$ when $t = \frac{2}{5}$: $0.3 = U \times \frac{2}{5} \times e^{-1}$

$U = \frac{3}{4}e = 2.0387\ldots \therefore$ speed $= 2.04\ m\ s^{-1}$

6 (d) Critical damping, because the auxiliary equation has a repeated root ($b^2 - 4ac = 20^2 - 4(4)(25) = 0$).