GCE A LEVEL MARKING SCHEME

SUMMER 2025

A LEVEL
FURTHER MATHEMATICS
UNIT 4 FURTHER PURE MATHEMATICS B
1305U40-1

© WJEC CBAC Ltd.

About this marking scheme

The purpose of this marking scheme is to provide teachers, learners, and other interested parties, with an understanding of the assessment criteria used to assess this specific assessment.

This marking scheme reflects the criteria by which this assessment was marked in a live series and was finalised following detailed discussion at an examiners' conference. A team of qualified examiners were trained specifically in the application of this marking scheme. The aim of the conference was to ensure that the marking scheme was interpreted and applied in the same way by all examiners. It may not be possible, or appropriate, to capture every variation that a candidate may present in their responses within this marking scheme. However, during the training conference, examiners were guided in using their professional judgement to credit alternative valid responses as instructed by the document, and through reviewing exemplar responses.

Without the benefit of participation in the examiners' conference, teachers, learners and other users, may have different views on certain matters of detail or interpretation. Therefore, it is strongly recommended that this marking scheme is used alongside other guidance, such as published exemplar materials or Guidance for Teaching. This marking scheme is final and will not be changed, unless in the event that a clear error is identified, as it reflects the criteria used to assess candidate responses during the live series.

© WJEC CBAC Ltd

WJEC GCE A LEVEL FURTHER MATHEMATICS

UNIT 4 FURTHER PURE MATHEMATICS B

SUMMER 2025 MARK SCHEME

Q	Solution	Mark	Notes
1. a)	Use of formula booklet expansion for e^x with x^2: $e^{x^2} = 1 + (x^2) + \frac{(x^2)^2}{2!} + \frac{(x^2)^3}{3!} + \cdots$ $e^{x^2} = 1 + x^2 + \frac{x^4}{2} + \frac{x^6}{6} + \cdots$	M1 A1 [2]	
b)	Approximate $\int_0^1 e^{x^2}\,dx = \int_0^1 \left(1 + x^2 + \frac{x^4}{2} + \frac{x^6}{6}\right) dx$ $= \left[x + \frac{x^3}{3} + \frac{x^5}{10} + \frac{x^7}{42}\right]_0^1$ $= \left(1 + \frac{1}{3} + \frac{1}{10} + \frac{1}{42}\right) - 0$ $\left(= \frac{51}{35}\right) = 1.457$ correct to 3dp	M1 A1 A1 [3]	FT (a), attempt to integrate cao Final A0 for exact answer 1.463 to 3dp
		[5]	

© WJEC CBAC Ltd

Q	Solution	Mark	Notes
2. a)	The result is true for $n = 1$ since it gives		
	$(\cos\theta + \mathrm{i}\sin\theta)^1 = \cos 1\theta + \mathrm{i}\sin 1\theta.$	B1	
	Let the results be true for $n = k$, ie	M1	
	$(\cos\theta + \mathrm{i}\sin\theta)^k = \cos k\theta + \mathrm{i}\sin k\theta$		
	Consider $n = k + 1$		
	$(\cos\theta + \mathrm{i}\sin\theta)^{k+1} = (\cos\theta + \mathrm{i}\sin\theta)^k(\cos\theta + \mathrm{i}\sin\theta)$	M1	
	$= (\cos k\theta + \mathrm{i}\sin k\theta)(\cos\theta + \mathrm{i}\sin\theta)$	A1	
	$= \cos k\theta\cos\theta - \sin k\theta\sin\theta + \mathrm{i}(\sin k\theta\cos\theta + \cos k\theta\sin\theta)$	A1	
	$= \cos(k+1)\theta + \mathrm{i}\sin(k+1)\theta$	A1	
	True for $n = k \Rightarrow$ true for $n = k + 1$ and since true for $n = 1$		
	the result is proved by mathematical induction.	A1	cso
		[7]	
b)	METHOD 1:		
	$(\cos\theta + \mathrm{i}\sin\theta)^4$		
	$= \cos^4\theta + 4\mathrm{i}\cos^3\theta\sin\theta + 6\mathrm{i}^2\cos^2\theta\sin^2\theta + 4\mathrm{i}^3\cos\theta\sin^3\theta + \mathrm{i}^4\sin^4\theta$	M1	Accept real terms only
	$= \cos^4\theta + 4\mathrm{i}\cos^3\theta\sin\theta - 6\cos^2\theta\sin^2\theta - 4\mathrm{i}\cos\theta\sin^3\theta + \sin^4\theta$	A1	Accept real terms only
	$(\cos\theta + \mathrm{i}\sin\theta)^4 = \cos 4\theta + \mathrm{i}\sin 4\theta$	B1	
	Equating real parts,		
	$\therefore \cos 4\theta = \cos^4\theta - 6\cos^2\theta\sin^2\theta + \sin^4\theta$		
	$= (1 - \sin^2\theta)^2 - 6(1 - \sin^2\theta)\sin^2\theta + \sin^4\theta$	M1	Use of identity
	$= 1 - 2\sin^2\theta + \sin^4\theta - 6\sin^2\theta + 6\sin^4\theta + \sin^4\theta$	A1	
	$= 8\sin^4\theta - 8\sin^2\theta + 1$	A1	Convincing
	METHOD 2:		
	Use of identity $\cos 4\theta = 1 - 2\sin^2 2\theta$	(M1)	
	Use of identity $\sin 2\theta = 2\sin\theta\cos\theta$	(M1)	
	$\therefore \cos 4\theta = 1 - 2(2\sin\theta\cos\theta)^2$	(A1)	
	$= 1 - 8\sin^2\theta\cos^2\theta$	(A1)	
	$= 1 - 8\sin^2\theta(1 - \sin^2\theta)$	(M1)	Use of identity
	$= 1 - 8\sin^2\theta + 8\sin^4\theta$	(A1)	Convincing
		[6]	

© WJEC CBAC Ltd

Q	Solution	Mark	Notes
2. c)i)	METHOD 1: $8\sin^4\theta - 8\sin^2\theta + 1 = 2\sin^2\theta - 1$ $8\sin^4\theta - 10\sin^2\theta + 2 = 0$ $4\sin^4\theta - 5\sin^2\theta + 1 = 0$ $(4\sin^2\theta - 1)(\sin^2\theta - 1) = 0$	M1	Accept degrees or radian throughout Solvable form
	Solving, $\sin\theta = \pm\frac{1}{2}$ or $\sin\theta = \pm 1$	m1 A2	A1 for 2 solutions
	$\theta = 30°, 150°, 210°, 330°\ldots$ or $\theta = 90°, 270°\ldots$		Accept $\pm\theta$ values
	$\theta = 360n° + 30°$, $\theta = 360n° + 150°$, $\theta = 360n° + 210°$, $\theta = 360n° + 330°$ $\theta = 360n° + 90°$, $\theta = 360n° + 270°$	A1 A1	For all 4 solns, oe For both solns, oe
	METHOD 2: $\cos 4\theta = 2\sin^2\theta - 1 = -\cos 2\theta$ $\cos 4\theta + \cos 2\theta = 0$	(B1)	Identifying identity
	Use of summation formulae, $2\cos\frac{4\theta+2\theta}{2}\cos\frac{4\theta-2\theta}{2} = 0$ $2\cos 3\theta\cos\theta = 0$	(M1) (A1)	
	Therefore, $\cos 3\theta = 0$ or $\cos\theta = 0$ $3\theta = 180n° + 90°$ $\theta = 60n° + 30°$ $\quad\quad$ $\theta = 180n° + 90°$	(m1) (A2)	Either A1 each set
ii)	Therefore, the general solution is $\theta = 60n° + 30°$	B1	oe
		[7]	
	Radian values: <table><tr><td>$30°$</td><td>$150°$</td><td>$210°$</td><td>$330°$</td></tr><tr><td>$\frac{\pi}{6}$</td><td>$\frac{5\pi}{6}$</td><td>$\frac{7\pi}{6}$</td><td>$\frac{11\pi}{6}$</td></tr><tr><td>$90°$</td><td>$270°$</td><td>$180°$</td><td>$360°$</td></tr><tr><td>$\frac{\pi}{2}$</td><td>$\frac{3\pi}{2}$</td><td>π</td><td>2π</td></tr><tr><td colspan="2">$60n° + 30°$</td><td colspan="2">$\frac{n\pi}{3} + \frac{\pi}{6}$</td></tr></table>		
		[20]	

© WJEC CBAC Ltd

Q	Solution	Mark	Notes
3. a)	$\cosh x^2 = \dfrac{e^{x^2} + e^{-x^2}}{2}$		
	$\dfrac{d}{dx}(\cosh x^2)$	M1	Differentiating
	$= \dfrac{2xe^{x^2} - 2xe^{-x^2}}{2}$	A1	
	$= 2x\left(\dfrac{e^{x^2} - e^{-x^2}}{2}\right)$		
	$= 2x \sinh x^2$	A1	Convincing, answer given
		[3]	
b)	METHOD 1: $\int 2x^3 \sinh x^2 \, dx$		
	Integrating by parts with $u = x^2, \frac{dv}{dx} = 2x \sinh x^2$	M1 m1	
	$= x^2 \cosh x^2 - \int 2x \cosh x^2 \, dx$	A1	
	$= x^2 \cosh x^2 - \sinh x^2 + c$	A1	ISW
	METHOD 2: $\int 2x^3 \sinh x^2 \, dx$		
	Integration by substitution with $u = x^2$ $du = 2x dx$	(M1)	
	leading to $\int u \sinh u \, du$	(A1)	
	Integration by parts with $u = u, \frac{dv}{dx} = \sinh u$	(m1)	
	$= u \cosh u - \int \cosh u \, du$		
	$= u \cosh u - \sinh u + c$		
	$= x^2 \cosh x^2 - \sinh x^2 + c$	(A1)	ISW
		[4]	
		[7]	

© WJEC CBAC Ltd

Q	Solution	Mark	Notes
4. a)	$\det\mathbf{M} = (3 \times 14) + (4 \times -4) = 26$	B1	
	Cofactor matrix: $\begin{pmatrix} 14 & -23 & -4 \\ 0 & 13 & 0 \\ -8 & 15 & 6 \end{pmatrix}$	B2	si B1 for any 5 correct
	Adjugate matrix: $\begin{pmatrix} 14 & 0 & -8 \\ -23 & 13 & 15 \\ -4 & 0 & 6 \end{pmatrix}$	B1	FT cofactor from B1
	$\therefore \mathbf{M}^{-1} = \frac{1}{26}\begin{pmatrix} 14 & 0 & -8 \\ -23 & 13 & 15 \\ -4 & 0 & 6 \end{pmatrix}$	B1 [5]	cao No workings, no marks
b)	Multiplying matrices and use of matrix equation, leading to $p + 4r + 2p = -6 \qquad (3p + 4r = -6)$ $3p - r + 2q = 11 \qquad (3p + 2q - r = 11)$ $2p + 5r + 2r = -17 \qquad (2p + 7r = -17)$ THEN:	M1 A2	A1 for 2 correct equations
	METHOD 1: $\begin{pmatrix} 3 & 0 & 4 \\ 3 & 2 & -1 \\ 2 & 0 & 7 \end{pmatrix}\begin{pmatrix} p \\ q \\ r \end{pmatrix} = \begin{pmatrix} -6 \\ 11 \\ -17 \end{pmatrix}$ Use of inverse from part (a): $\begin{pmatrix} p \\ q \\ r \end{pmatrix} = \frac{1}{26}\begin{pmatrix} 14 & 0 & -8 \\ -23 & 13 & 15 \\ -4 & 0 & 6 \end{pmatrix}\begin{pmatrix} -6 \\ 11 \\ -17 \end{pmatrix}$ $\begin{pmatrix} p \\ q \\ r \end{pmatrix} = \begin{pmatrix} 2 \\ 1 \\ -3 \end{pmatrix}$	M1 A1 A2	FT (a) cao A1 for 2 correct solutions
	METHOD 2: $\left(\begin{array}{ccc\|c} 3 & 0 & 4 & -6 \\ 3 & 2 & -1 & 11 \\ 2 & 0 & 7 & -17 \end{array}\right)$ Using row reduction e.g. r1 – 1.5r3 and r2 – 1.5r3: $\left(\begin{array}{ccc\|c} 0 & 0 & -6.5 & 19.5 \\ 0 & 2 & -11.5 & 36.5 \\ 2 & 0 & 7 & -17 \end{array}\right)$ Therefore, $-6.5r = 19.5 \qquad \therefore r = -3$ $2q + (-11.5 \times -3) = 36.5 \qquad \therefore q = 1$ $2p + (7 \times -3) = -17 \qquad \therefore p = 2$	(M1) (A1) (A2)	cao A1 for 2 correct solutions
	METHOD 3: Using 1st and 3rd equations, removal of 1 variable $p = 2$ or $r = -3$ Other variable Use of second equation to find q: $(3 \times 2) + 2q - (-3) = 11 \qquad \therefore q = 1$	(M1) (A1) (A1) (A1) [7]	cao cao
		[12]	

© WJEC CBAC Ltd

Q	Solution	Mark	Notes
5.	METHOD 1:		
	Volume = $\pi \int_0^1 \sinh^2 2y \, \mathrm{d}y$	B1	Correct notation required
	Use of $\cosh 4y = 1 + 2\sinh^2 2y$	M1	M0 no workings
	$\pi \int_0^1 \left(\frac{\cosh 4y - 1}{2}\right) \mathrm{d}y$	A1	Integrable form with no more than 1 slip
	$\pi \left[\frac{1}{2} \times \frac{\sinh 4y}{4} - \frac{1}{2}y\right]_0^1$	A1	oe cao integration
	$\pi \left[\left(3.411 \ldots - \frac{1}{2}\right) - (0)\right]$	m1	Substitute in correct limits
	Volume = 9.146	A1	cao
	METHOD 2:		
	Volume = $\pi \int_0^1 \sinh^2 2y \, \mathrm{d}y$	(B1)	Correct notation required
	Use of $\sinh 2y = \frac{\mathrm{e}^{2y} - \mathrm{e}^{-2y}}{2}$	(M1)	M0 no workings
	$\frac{\pi}{4} \int_0^1 (\mathrm{e}^{4y} - 2 + \mathrm{e}^{-4y}) \mathrm{d}y$	(A1)	Integrable form with no more than 1 slip
	$\frac{\pi}{4} \left[\frac{\mathrm{e}^{4y}}{4} - 2y - \frac{\mathrm{e}^{-4y}}{4}\right]_0^1$	(A1)	oe cao integration
	$\frac{\pi}{4} \left[\left(\frac{\mathrm{e}^4}{4} - 2 - \frac{\mathrm{e}^{-4}}{4}\right) - \left(\frac{1}{4} - 0 - \frac{1}{4}\right)\right]$	(m1)	Substitute in correct limits
	Volume = 9.146	(A1)	cao
		[6]	If M0, award SC1 for 9.1... unsupported
		[6]	

© WJEC CBAC Ltd

Q	Solution	Mark	Notes
6.	Dividing both sides by $\cos x$: $\frac{dy}{dx} + \frac{y}{\cos x} = \tan x$	M1	
	Integrating factor: $e^{\int \frac{1}{\cos x} dx} = e^{\int \sec x dx}$ $= e^{\ln(\sec x + \tan x)} = \sec x + \tan x$	M1 A1	
	Multiplying both sides: $(\sec x + \tan x)\frac{dy}{dx} + y\frac{(\sec x + \tan x)}{\cos x} = (\sec x + \tan x)\tan x$	M1	
	RHS: $\sec x \tan x + \tan^2 x \rightarrow \sec x \tan x + \sec^2 x - 1$	A1	Right-side integrable form
	Integrating both sides: $y(\sec x + \tan x) = \sec x + \tan x - x + c$	m1 A1	
	Substituting $3 = 1 + 0 - 0 + c \ \rightarrow \ c = 2$		
	Solution: $y(\sec x + \tan x) = \sec x + \tan x - x + 2$ $y = \frac{\sec x + \tan x - x + 2}{\sec x + \tan x}$	B1	oe
		[8]	
		[8]	

© WJEC CBAC Ltd

Q	Solution	Mark	Notes
7. a) i)	METHOD 1: $\frac{dy}{dx} = -\frac{1}{\sqrt{1-(2x-1)^2}} \times 2$ $\frac{dy}{dx} = -\frac{2}{\sqrt{1-4x^2+4x-1}}$ $\frac{dy}{dx} = -\frac{1}{\sqrt{x-x^2}}$	M1 A1 A1	Must be chain rule oe oe simplified
	METHOD 2: $\cos y = 2x - 1$ $-\sin y \frac{dy}{dx} = 2$ $\frac{dy}{dx} = -\frac{2}{\sin y}$ $\frac{dy}{dx} = -\frac{2}{\sqrt{1-\cos^2 y}}$ $\frac{dy}{dx} = -\frac{2}{\sqrt{1-(2x-1)^2}}$ $\frac{dy}{dx} = -\frac{2}{\sqrt{1-4x^2+4x-1}}$ $\frac{dy}{dx} = -\frac{1}{\sqrt{x-x^2}}$	(M1) (A1) (A1) [3]	 oe oe simplified
ii)	$\frac{dy}{dx} = (3x \times \cosh 4x \times 4) + (\sinh 4x \times 3)$ $\frac{dy}{dx} = 12x \cosh 4x + 3 \sinh 4x$ or $3(4x \cosh 4x + \sinh 4x)$	M1 A1 A1 [3]	Product rule One term correct Fully correct and simplified
b)	$\frac{dy}{dx} = \frac{(\cosh x \times -\sin x) - (\cos x \times \sinh x)}{(\cosh x)^2}$ At $x = \frac{\pi}{2}$, $\frac{dy}{dx} = -\frac{1}{\cosh\frac{\pi}{2}} = -0.3985\ldots$ and $y = 0$ Gradient of normal: $\cosh\frac{\pi}{2} = \frac{-1}{-0.3985\ldots} = 2.509\ldots$ Equation: $y - 0 = 2.509\ldots\left(x - \frac{\pi}{2}\right)$ $y = 2.51x - 3.94$	M1 A1 B1 B1 m1 A1 [6]	Quotient rule oe Fully correct Both dy/dx and y FT their dy/dx Dependant on M1 Must be 2d.p.
		[12]	

© WJEC CBAC Ltd

Q	Solution	Mark	Notes
8. a)	$4\cos\theta + 7\sin\theta = 4 - \tan\left(\frac{\theta}{2}\right)$ $4\left(\frac{1-t^2}{1+t^2}\right) + 7\left(\frac{2t}{1+t^2}\right) = 4 - t$ $4(1-t^2) + 7(2t) = (4-t)(1+t^2)$ $4 - 4t^2 + 14t = 4 + 4t^2 - t - t^3$ $t^3 - 8t^2 + 15t = 0$	M1 A1 A1 [3]	Use of t formulae Removing fraction Convincing
b)	$t(t^2 - 8t + 15) = 0$ Solving, $t = 0$ or $t^2 - 8t + 15 = 0$ $(t-3)(t-5) = 0$ $t = 0$ $t = 3$ or $t = 5$ $\tan\left(\frac{\theta}{2}\right) = 0$ $\tan\left(\frac{\theta}{2}\right) = 3$ or $\tan\left(\frac{\theta}{2}\right) = 5$ $\theta = 0$ $\theta = 2.498\ldots$ $\theta = 2.7468\ldots$ *Other factorisations could include (not exclusively)* • $(t-3)(t^2-5t)$ • $(t-5)(t^2-3t)$ • *Completed squares*	B1 M1 A2 B2 [6]	si M0 no workings A1 for any two B1 for any two Accept degrees Ignore angles outside range -1 for each additional roots in range
		[9]	

© WJEC CBAC Ltd

Q	Solution	Mark	Notes
9.	Solve auxiliary $m^2 + 16 = 0$ $m^2 = -16$ $m = \pm 4\text{i}$	M1	
	Complementary function: $y = A\sin 4x + B\cos 4x$	A1	
	Use particular integral of the form $C\sin x + D\cos x$ $\frac{\mathrm{d}y}{\mathrm{d}x} = C\cos x - D\sin x$ $\frac{\mathrm{d}^2y}{\mathrm{d}x^2} = -C\sin x - D\cos x$	M1 A1	
	Therefore, $-C\sin x - D\cos x + 16(C\sin x + D\cos x) = 3\sin x$	A1	Substitution
	$-C + 16C = 3 \quad \rightarrow \quad C = \frac{1}{5}$ $-D + 16D = 0 \quad \rightarrow \quad D = 0$	A1	Both values *FT C,D for M1A1M1A1*
	General Solution: $y = A\sin 4x + B\cos 4x + \frac{1}{5}\sin x$	M1	Sub and differentiate
	$\frac{\mathrm{d}y}{\mathrm{d}x} = 4A\cos 4x - 4B\sin 4x + \frac{1}{5}\cos x$	A1	
	When $x = \pi$, $y = 3$, $\frac{\mathrm{d}y}{\mathrm{d}x} = 0$ $B = 3$ and $\frac{\mathrm{d}y}{\mathrm{d}x} = 4A - \frac{1}{5} = 0 \rightarrow \quad A = \frac{1}{20}$	M1 A1	Substituting π and one other value into either y or $\frac{\mathrm{d}y}{\mathrm{d}x}$ Both A and B
	Therefore, $y = \frac{1}{20}\sin 4x + 3\cos 4x + \frac{1}{5}\sin x$	B1	cao
		[11]	
		[11]	

© WJEC CBAC Ltd

Q	Solution	Mark	Notes
10. a)i)	$\frac{2x^3+x^2+12}{x^3-2x^2-3x+6} = \frac{2(x^3-2x^2-3x+6)}{x^3-2x^2-3x+6} + \frac{5x^2+6x}{x^3-2x^2-3x+6}$ $\therefore \frac{2x^3+x^2+12}{x^3-2x^2-3x+6} = 2 + \frac{5x^2+6x}{x^3-2x^2-3x+6}$	M1 A1 [2]	Or polynomial division convincing
ii)	METHOD 1: $(x-2)$ is a linear factor of x^3-2x^2-3x+6 $\therefore \frac{2x^3+x^2+12}{x^3-2x^2-3x+6} = 2 + \frac{5x^2+6x}{(x-2)(x^2-3)}$ $\frac{5x^2+6x}{(x-2)(x^2-3)} = \frac{A}{x-2} + \frac{Bx+C}{x^2-3}$ $5x^2+6x = A(x^2-3) + (Bx+C)(x-2)$ When $x=2$, $32 = A$ When $x=0$, $0 = -3A - 2C \quad \rightarrow C = -48$ Comparing x^2 coefficients: $5 = A + B \quad \rightarrow B = -27$ Therefore, $\int \left(\frac{2x^3+x^2+12}{x^3-2x^2-3x+6}\right) \mathrm{d}x = \int \left(2 + \frac{32}{x-2} + \frac{-27x-48}{x^2-3}\right) \mathrm{d}x$ $= \int \left(2 + \frac{32}{x-2} + \frac{-27x}{x^2-3} + \frac{-48}{x^2-3}\right) \mathrm{d}x$ $= 2x + 32\ln\lvert x-2\rvert - \frac{27}{2}\ln\lvert x^2-3\rvert - \frac{48}{2\sqrt{3}}\ln\left\lvert\frac{x-\sqrt{3}}{x+\sqrt{3}}\right\rvert + c$ oe	B1 M1 m1 A2 m1 A3	si Partial fractions All values A1 two values Integration FT their partial fractions Fully correct ISW A2 for 4 correct terms A1 for 2 correct terms Note: $\frac{48}{2\sqrt{3}} = \frac{24}{\sqrt{3}} = 8\sqrt{3}$


© WJEC CBAC Ltd

Q	Solution	Mark	Notes
10. a) ii)	METHOD 2: $(x-2)$ is a linear factor of x^3-2x^2-3x+6	(B1)	si
	$\therefore \frac{2x^3+x^2+12}{x^3-2x^2-3x+6} = 2+\frac{5x^2+6x}{(x-2)(x^2-3)}$		
	$\frac{5x^2+6x}{(x-2)(x^2-3)} = \frac{A}{x-2}+\frac{B}{x+\sqrt{3}}+\frac{C}{x-\sqrt{3}}$	(M1)	Partial fractions
	$5x^2+6x = A(x+\sqrt{3})(x-\sqrt{3})+B(x-\sqrt{3})(x-2)+C(x+\sqrt{3})(x-2)$	(m1)	
	When $x=2$, $\quad 32=A$ When $x=\sqrt{3}$, $\quad 15+6\sqrt{3}=C(\sqrt{3}+\sqrt{3})(\sqrt{3}-2)$ $\frac{15+6\sqrt{3}}{6-4\sqrt{3}} = C \quad \left(=\frac{-16\sqrt{3}-27}{2}\right)$ When $x=-\sqrt{3}$, $15-6\sqrt{3}=B(-\sqrt{3}-\sqrt{3})(-\sqrt{3}-2)$ $\frac{15-6\sqrt{3}}{6+4\sqrt{3}} = B \quad \left(=\frac{16\sqrt{3}-27}{2}\right)$	(A2)	All values A1 two values
	Therefore, $\int\left(\frac{2x^3+x^2+12}{x^3-2x^2-3x+6}\right)\mathrm{d}x = \int\left(2+\frac{32}{x-2}+\frac{\frac{15-6\sqrt{3}}{6+4\sqrt{3}}}{x+\sqrt{3}}+\frac{\frac{15+6\sqrt{3}}{6-4\sqrt{3}}}{x-\sqrt{3}}\right)\mathrm{d}x$	(m1)	Integration FT their partial fractions
	$= 2x+32\ln\lvert x-2\rvert+\frac{15-6\sqrt{3}}{6+4\sqrt{3}}\ln\lvert x+\sqrt{3}\rvert+\frac{15+6\sqrt{3}}{6-4\sqrt{3}}\ln\lvert x-\sqrt{3}\rvert+c$ oe e.g. with rationalised coefficients: $= 2x+32\ln\lvert x-2\rvert+\frac{16\sqrt{3}-27}{2}\ln\lvert x+\sqrt{3}\rvert+\frac{-16\sqrt{3}-27}{2}\ln\lvert x-\sqrt{3}\rvert+c$	(A3)	Fully correct ISW A2 for 4 correct terms A1 for 2 correct terms
		[9]	

© WJEC CBAC Ltd

Q	Solution	Mark	Notes
10. b)	$4-4x-x^2 = 8-(x+2)^2$	B1	
	$\int_{-2}^{0} \frac{1}{\sqrt{4-4x-x^2}}\,\mathrm{d}x = \int_{-2}^{0} \frac{1}{\sqrt{8-(x+2)^2}}\,\mathrm{d}x$	M1	FT square for $a^2 - x^2$ only
	$= \left[\sin^{-1}\left(\frac{x+2}{\sqrt{8}}\right)\right]_{-2}^{0}$	A1	
	$= \frac{\pi}{4} - 0$		
	$= \frac{\pi}{4}$	A1	cao No workings, no marks
		[4]	
c)	METHOD 1: Mean value $= \frac{1}{2-0}\int_0^2 \tanh 2x\,\mathrm{d}x$	M1	Attempt to integrate. Condone omission of ½ for M1A1
	$= \frac{1}{2}\left[\frac{\ln(\cosh 2x)}{2}\right]_0^2$	A1	Integration
	$= \frac{1}{2}(1.65359\ldots - 0)$		
	$= 0.83$	A1	cao
	METHOD 2: $\tanh 2x = \frac{\mathrm{e}^{2x}-\mathrm{e}^{-2x}}{\mathrm{e}^{2x}+\mathrm{e}^{-2x}}$		
	Mean value $= \frac{1}{2-0}\int_0^2 \tanh 2x\,\mathrm{d}x$ $= \frac{1}{2}\int_0^2 \frac{\mathrm{e}^{2x}-\mathrm{e}^{-2x}}{\mathrm{e}^{2x}+\mathrm{e}^{-2x}}\,\mathrm{d}x$	M1	Attempt to integrate. Condone omission of ½ for M1A1
	Integration by substitution with $u = \mathrm{e}^{2x} + \mathrm{e}^{-2x}$ $\mathrm{d}u = (2\mathrm{e}^{2x} - 2\mathrm{e}^{-2x})\mathrm{d}x$ leading to $\frac{1}{2}\int \frac{\mathrm{d}u}{u}$ $= \frac{1}{2}\ln u$	A1	Integration
	With limits and $\frac{1}{2}$: $= \frac{1}{4}[\ln u]_{2e}^{\mathrm{e}^4 + \mathrm{e}^{-4}}$ OR $= \frac{1}{4}[\ln(\mathrm{e}^{2x} + \mathrm{e}^{-2x})]_0^2$		
	$= \frac{1}{4}(4.000335\ldots - 0.69314\ldots)$		
	$= 0.83$	A1	cao If M0, SC1 for correct answer only
		[3]	
		[18]	

© WJEC CBAC Ltd

Q	Solution	Mark	Notes
11. a)		G2 [2]	Fully correct graph Allow G1 for • two circles • half graph
b)	$y = r\sin\theta = \cos^2\theta\sin\theta$	M1	
	Differentiating $\frac{dy}{d\theta} = \cos^3\theta - 2\cos\theta\sin^2\theta$	m1 A1	Must be product rule
	When parallel to initial line $\frac{dy}{d\theta} = 0$		
	$\cos^3\theta - 2\cos\theta\sin^2\theta = 0$ $\cos\theta(\cos^2\theta - 2\sin^2\theta) = 0$	m1	
	$\cos\theta = 0$ (no tangent parallel to initial line) or $\cos^2\theta - 2\sin^2\theta = 0$ $\cos^2\theta - 2(1-\cos^2\theta) = 0$ $3\cos^2\theta - 2 = 0$	A1	
	$\cos\theta = \pm\sqrt{\frac{2}{3}}$ $(\sin\theta = \pm\frac{1}{\sqrt{3}}, \tan\theta = \pm\frac{1}{\sqrt{2}})$	A1	Any value
	$r = \frac{2}{3}$	A1	
	Then, $x = r\cos\theta = \frac{2}{3}\times\sqrt{\frac{2}{3}} = \frac{2\sqrt{2}}{3\sqrt{3}} = \frac{2\sqrt{6}}{9}$ $y = r\sin\theta = \frac{2}{3}\times\frac{1}{\sqrt{3}} = \frac{2}{3\sqrt{3}} = \frac{2\sqrt{3}}{9}$	m1	Attempt to find x, y for any point
	Area = $2x \times 2y$	m1	
	$= 2\times\frac{2\sqrt{2}}{3\sqrt{3}}\times 2\times\frac{2}{3\sqrt{3}} = \frac{16\sqrt{2}}{27}$ oe	A1 [10]	Must be exact
	Alternative possible derivatives: • $y = \left(\frac{\cos 2\theta+1}{2}\right)\sin\theta$ $\frac{dy}{d\theta} = \left(\frac{\cos 2\theta+1}{2}\right)\cos\theta - \sin\theta\sin 2\theta$ • $y = \left(\frac{\cos 2\theta+1}{2}\right)\sin\theta = \frac{1}{2}\cos 2\theta\sin\theta + \frac{1}{2}\sin\theta$ $\frac{dy}{d\theta} = \frac{1}{2}\cos 2\theta\sin\theta - \sin 2\theta\sin\theta + \frac{1}{2}\cos\theta$ • $y = (1-\sin^2\theta)\sin\theta = \sin\theta - \sin^3\theta$ $\frac{dy}{d\theta} = \cos\theta - 3\cos\theta\sin^2\theta$		
		[12]	

© WJEC CBAC Ltd